abcam

Product datasheet

HRP Anti-muscle Actin antibody [EPR8484] - Loading Control ab185058

יעלאעבע RabMAb

★★★★★ 1 Abreviews 2 References

画像数 2

製品の概要

製品名 HRP Anti-muscle Actin antibody [EPR8484] - Loading Control

製品の詳細 HRP Rabbit monoclonal [EPR8484] to muscle Actin - Loading Control

由来種 Rabbit 標識 HRP

アプリケーション **適用あり:** WB

種交差性 交差種: Mouse, Rat, Human

免疫原 Synthetic peptide within Human muscle Actin aa 350-450. The exact immunogen sequence used

> to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific

Support team to discuss your requirements.

Database link: P68032

Run BLAST with Run BLAST with

ポジティブ・コントロール NIH 3T3 cell lysate; Human fetal artery, kidney and heart lisates; Human uterus, stomach and skeletal muscle lysates; Human skeletal muscle tissue; Human smooth muscle in colon tissue;

Human heart muscle tissue.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® patents.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

1

バッファー pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), PBS, 1% BSA

精製度 Protein A purified

ポリ/モノ モノクローナル **クローン名** EPR8484

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab185058の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★☆ (1)	1/10000. Detects a band of approximately 42 kDa (predicted molecular weight: 42 kDa).

ターゲット情報

機能

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

関連疾患

Defects in ACTA1 are the cause of nemaline myopathy type 3 (NEM3) [MIM:161800]. A form of nemaline myopathy. Nemaline myopathies are muscular disorders characterized by muscle weakness of varying severity and onset, and abnormal thread-or rod-like structures in muscle fibers on histologic examination. The phenotype at histological level is variable. Some patients present areas devoid of oxidative activity containg (cores) within myofibers. Core lesions are unstructured and poorly circumscribed.

Defects in ACTA1 are a cause of myopathy, actin, congenital, with excess of thin myofilaments (MPCETM) [MIM:161800]. A congenital muscular disorder characterized at histological level by areas of sarcoplasm devoid of normal myofibrils and mitochondria, and replaced with dense masses of thin filaments. Central cores, rods, ragged red fibers, and necrosis are absent. Defects in ACTA1 are a cause of congenital myopathy with fiber-type disproportion (CFTD) [MIM:255310]; also known as congenital fiber-type disproportion myopathy (CFTDM). CFTD is a genetically heterogeneous disorder in which there is relative hypotrophy of type 1 muscle fibers compared to type 2 fibers on skeletal muscle biopsy. However, these findings are not specific and can be found in many different myopathic and neuropathic conditions.


配列類似性 Belongs to the actin family.

翻訳後修飾 Oxidation of Met-46 by MICALs (MICAL1, MICAL2 or MICAL3) to form methionine sulfoxide promotes actin filament depolymerization. Methionine sulfoxide is produced stereospecifically, but

it is not known whether the (S)-S-oxide or the (R)-S-oxide is produced.

細胞内局在 Cytoplasm > cytoskeleton.

画像



Western blot - HRP Anti-muscle Actin antibody
[EPR8484] - Loading Control (ab185058)

All lanes : HRP Anti-muscle Actin antibody [EPR8484] - Loading Control (ab185058) at 1/10000 dilution

Lane 1 : NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate at 20 μg

Lane 2 : Human kidney tissue lysate - total protein (ab30203) at 20 µq

Lane 3: Human Fetal Heart at 5 µg

Lane 4: Human skeletal muscle tissue lysate - total protein

(ab29330) at 10 µg

Lane 5: Skeletal Muscle (Rat) Tissue Lysate at 10 µg

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 42 kDa

Additional bands at: 42 kDa. We are unsure as to the identity of

these extra bands.

Exposure time: 5 seconds

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab185058 overnight at 4°C. Antibody binding was visualised using ECL development solution ab133406







HRP Anti-muscle Actin antibody [EPR8484] -Loading Control (ab185058)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors