

HRP Anti-Mycobacterium tuberculosis antibody ab21189

医薬用外劇物

製品の概要

製品名	HRP Anti-Mycobacterium tuberculosis antibody
製品の詳細	HRP Rabbit polyclonal to Mycobacterium tuberculosis
由来種	Rabbit
標識	HRP
アプリケーション	適用あり: IHC-P
種交差性	交差種: Mycobacterium tuberculosis
免疫原	Tissue, cells or virus corresponding to Mycobacterium tuberculosis. Mycobacterium tuberculosis purified protein derivative
特記事項	<p>IgG fraction covalently coupled to a highly purified preparation of horseradish peroxidase (RZ>3). Care is taken to ensure adequate conjugation while reserving maximum enzyme activity. Free enzyme is absent. Estimated Molar HRP: IgG substitution is 2-3</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	Preservative: 0.002% Thimerosal (merthiolate) Constituents: PBS, 1% BSA
精製度	IgG fraction
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab21189の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent dilution.

ターゲット情報

関連性 Mycobacterium tuberculosis is the most common cause of tuberculosis. Primary infection begins with inhalation of 1 to 10 aerosolised bacilli. The pathogenicity of the organism is determined by its ability to escape host immune responses as well as eliciting delayed hypersensitivity. Alveolar macrophages engulf the invading cells but are unable to mount an effective defense. Several virulence factors are responsible for this apparent failure; most notably in the mycobacterial cell wall are the cord factor, lipoarabinomannan, and the 65 kd heat shock protein or HSP65. The emergence of new strains of resistant Mycobacterium tuberculosis has created new interest in clinical diagnosis. Studies have shown immunohistochemical techniques to be superior to conventional special stains. Thus the demonstration of mycobacterial antigens are not only useful in establishing mycobacterial aetiology, but can also be used as an alternative method to the conventional Ziehl-Neelsen method.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors