

# HRP Anti-O-Linked N-Acetylglucosamine antibody [RL2] ab201995

**2 References**   [画像数 2](#)

### 製品の概要

製品名	HRP Anti-O-Linked N-Acetylglucosamine antibody [RL2]
製品の詳細	HRP Mouse monoclonal [RL2] to O-Linked N-Acetylglucosamine
由来種	Mouse
標識	HRP
アプリケーション	<b>適用あり:</b> IHC-P, WB
種交差性	<b>交差種:</b> Human
ポジティブ・コントロール	IHC-P: Normal human colon tissue. WB: Jurkat treated with 0 uM PugNAc and Jurkat treated with 4 mM glucosamine and 50 uM PugNAc (3 hours).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Store In the Dark.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.1% Proclin 300 Solution</p> <p>Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA</p>
精製度	IgG fraction
ポリ/モノ	モノクローナル
クローン名	RL2
アイソタイプ	IgG1

## アプリケーション


**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab201995の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		1/1000.

## ターゲット情報

**関連性**      Many cellular proteins, including nuclear pore, oncogene, cytoskeletal, heat shock, viral and transcription regulatory proteins contain single O-linked N-acetylglucosamine (O-GlcNAc) residues attached to serine or threonine residues. It has been observed that O-GlcNAc glycosylated proteins tend to be under phosphorylated relative to unglycosylated proteins and that O-GlcNAc bearing proteins tend to be found in multimeric complexes. This has led to the suggestion that O-GlcNAc glycosylation may obscure phosphorylation sites and acts as a signaling mechanism or mediator of signaling.

## 画像



**All lanes :** HRP Anti-O-Linked N-Acetylglucosamine antibody [RL2] (ab201995) at 1/1000 dilution

**Lane 1 :** Jurkat treated with 0 uM PugNAc

**Lane 2 :** Jurkat treated with 4 mM glucosamine and 50 uM PugNAc (3 hours)

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Observed band size:** 100,65 kDa

**Exposure time:** 2 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab201995 overnight at 4°C. Antibody binding

was visualised using ECL development solution **ab133406**.



IHC image of O-linked N-Acetylglucosamine staining in a section of formalin-fixed paraffin-embedded normal human colon tissue\*, performed on a Leica BOND. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab201995, 1/100 dilution, for 15 mins at room temperature. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors