

Anti-Hsp90 beta antibody [E296] ab32568

KO 評価済 リコンビナント RabMAb

16 References 画像数 10

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-Hsp90 beta antibody [E296] |
| 製品の詳細 | Rabbit monoclonal [E296] to Hsp90 beta |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: WB, IHC-P, ICC/IF 適用なし: Flow Cyt |
| 種交差性 | 交差種: Mouse, Rat, Human |
| 免疫原 | Synthetic peptide within Human Hsp90 beta aa 1-100 (N terminal). The exact sequence is proprietary. |
| ポジティブ・コントロール | WB: Saos-2, HL-60, HEK293T, Jurkat, SH-SY5Y, Raji, A431 and HeLa whole cell lysate (ab150035); Mouse brain and heart tissue lysates; Rat brain and heart tissue lysates. IHC-P: Stomach and urinary bladder carcinoma tissues. ICC/IF: HepG2 cells. |
| 特記事項 | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

製品の特性

| | |
|-------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| バッファー | pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| 精製度 | Protein A purified |
| ポリ/モノ | モノクローナル |
| クローン名 | E296 |

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab32568の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| WB | | 1/100000 - 1/500000. Detects a band of approximately 92 kDa (predicted molecular weight: 83 kDa). For unpurified, use 1/500. |
| IHC-P | | 1/150. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> . |
| ICC/IF | | 1/100. |

追加情報

Is unsuitable for Flow Cyt.

ターゲット情報

機能

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function.

配列類似性

Belongs to the heat shock protein 90 family.

ドメイン

The TPR repeat-binding motif mediates interaction with TPR repeat-containing proteins.

翻訳後修飾

Ubiquitinated in the presence of STUB1-UBE2D1 complex (in vitro).


ISGylated.

S-nitrosylated; negatively regulates the ATPase activity.

細胞内局在

Cytoplasm. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

画像



Western blot - Anti-Hsp90 beta antibody [E296] (ab32568)

All lanes : Anti-Hsp90 beta antibody [E296] (ab32568) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : HSP90AB1 knockout HEK293T cell lysate

Lane 3 : Jurkat cell lysate

Lane 4 : SH-SY5Y cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) at 1/10000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa


Lanes 1-4: Merged signal (red and green). Green - ab32568 observed at 90 kDa. Red - loading control **ab8245** observed at 36 kDa.

ab32568 Anti-Hsp90 beta antibody [E296] was shown to specifically react with Hsp90 beta in wild-type HEK293T cells. Loss of signal was observed when knockout cell line **ab266117** (knockout cell lysate **ab257190**) was used. Wild-type and Hsp90 beta knockout samples were subjected to SDS-PAGE. ab32568 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated at room temperature for 2.5 hours at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp90 beta antibody [E296] (ab32568)

Immunohistochemical staining of paraffin embedded human stomach with purified ab32568 at a working dilution of 1 in 150. The secondary antibody used is a HRP polymer for rabbit IgG. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Immunocytochemistry/ Immunofluorescence - Anti-Hsp90 beta antibody [E296] (ab32568)

Immunofluorescence staining of HepG2 cells with purified ab32568 at a working dilution of 1 in 100, counter-stained with DAPI. The secondary antibody was Alexa Fluor[®] 555 goat anti rabbit (**ab150082**), used at a dilution of 1 in 400. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative control is shown in bottom right hand panel - for the negative control, purified ab32568 was used at a dilution of 1/200 followed by an Alexa Fluor[®] 488 goat anti-mouse antibody at a dilution of 1/500.



Western blot - Anti-Hsp90 beta antibody [E296]
(ab32568)

All lanes : Anti-Hsp90 beta antibody [E296] (ab32568) at 1/200000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : HSP90AB1 knockout HEK-293T cell lysate

Lane 3 : Saos-2 cell lysate

Lane 4 : HL-60 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 83 kDa

Observed band size: 85 kDa

Lanes 1 - 4: Merged signal (red and green). Green - ab32568 observed at 85 kDa. Red - loading control **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.

ab32568 was shown to react with Hsp90 beta in wild-type HEK-293T cells in western blot with loss of signal observed in HSP90AB1 knockout cell line **ab266117** (HSP90AB1 knockout cell lysate **ab257190**). Wild-type and HSP90AB1 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab32568 and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 200000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-Hsp90 beta antibody [E296] (ab32568) at 1/100000 dilution (purified)

Lane 1 : Mouse brain tissue lysate

Lane 2 : Mouse heart tissue lysate

Lane 3 : Rat brain tissue lysate

Lane 4 : Rat heart tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST



Western blot - Anti-Hsp90 beta antibody [E296]
(ab32568)


Anti-Hsp90 beta antibody [E296] (ab32568) at 1/100000 dilution
(purified) + SH-SH5Y cell lysate at 20 µg

Secondary

HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa



Western blot - Anti-Hsp90 beta antibody [E296]
(ab32568)

All lanes : Anti-Hsp90 beta antibody [E296] (ab32568) at
1/100000 dilution (purified)

Lane 1 : HeLa cell lysate

Lane 2 : Jurkat cell lysate

Lane 3 : Raji cell lysate

Lane 4 : A431 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST




Anti-Hsp90 beta antibody [E296] (ab32568) at 1/500 dilution
(unpurified) + Hela cell lysate

Predicted band size: 83 kDa

Observed band size: 92 kDa

Western blot - Anti-Hsp90 beta antibody [E296]
(ab32568)



Immunohistochemical analysis of paraffin-embedded human urinary bladder carcinoma using unpurified ab32568 at 1/50 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp90 beta antibody [E296] (ab32568)

Why choose a recombinant antibody?

| | |
|--|--|
|  <p>Research with confidence Consistent and reproducible results</p> |  <p>Long-term and scalable supply Recombinant technology</p> |
|  <p>Success from the first experiment Confirmed specificity</p> |  <p>Ethical standards compliant Animal-free production</p> |

Anti-Hsp90 beta antibody [E296] (ab32568)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors