


Anti-Ikaros antibody [EPR13790] - BSA and Azide free ab206645

KO 評価済 リコンビナント RabMAb

画像数 5

製品の概要

製品名	Anti-Ikaros antibody [EPR13790] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR13790] to Ikaros - BSA and Azide free
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), IHC-P, WB
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Jurkat, Daudi, Raji, Ramos and MOLT4 cell lysates; Human thymus and mouse spleen tissues; MOLT4 cells.
特記事項	<p>ab206645 is the carrier-free version of ab191394.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR13790
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab206645の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Detects a band of approximately 50-70 kDa (predicted molecular weight: 58 kDa).

ターゲット情報

機能	Transcription regulator of hematopoietic cell differentiation (PubMed:17934067). Binds gamma-satellite DNA (PubMed:17135265, PubMed:19141594). Plays a role in the development of lymphocytes, B- and T-cells. Binds and activates the enhancer (delta-A element) of the CD3-delta gene. Repressor of the TDT (fukuzterminal deoxynucleotidyltransferase) gene during thymocyte differentiation. Regulates transcription through association with both HDAC-dependent and HDAC-independent complexes. Targets the 2 chromatin-remodeling complexes, NuRD and BAF (SWI/SNF), in a single complex (PYR complex), to the beta-globin locus in adult erythrocytes. Increases normal apoptosis in adult erythroid cells. Confers early temporal competence to retinal progenitor cells (RPCs) (By similarity). Function is isoform-specific and is modulated by dominant-negative inactive isoforms (PubMed:17135265, PubMed:17934067).
組織特異性	Abundantly expressed in thymus, spleen and peripheral blood Leukocytes and lymph nodes. Lower expression in bone marrow and small intestine.
関連疾患	Defects in IKZF1 are frequent occurrences (28.6%) in acute lymphoblastic leukemia (ALL). Such alterations or deletions lead to poor prognosis for ALL. Chromosomal aberrations involving IKZF1 are a cause of B-cell non-Hodgkin lymphomas (B-cell

	NHL). Translocation t(3;7)(q27;p12), with BCL6.
配列類似性	Belongs to the Ikaros C2H2-type zinc-finger protein family. Contains 6 C2H2-type zinc fingers.
ドメイン	The N-terminal zinc-fingers 2 and 3 are required for DNA binding as well as for targeting IKZF1 to pericentromeric heterochromatin. The C-terminal zinc-finger domain is required for dimerization.
翻訳後修飾	Phosphorylation controls cell-cycle progression from late G(1) stage to S stage. Hyperphosphorylated during G2/M phase. Dephosphorylated state during late G(1) phase. Phosphorylation on Thr-140 is required for DNA and pericentromeric location during mitosis. CK2 is the main kinase, in vitro. GSK3 and CDK may also contribute to phosphorylation of the C-terminal serine and threonine residues. Phosphorylation on these C-terminal residues reduces the DNA-binding ability. Phosphorylation/dephosphorylation events on Ser-13 and Ser-295 regulate TDT expression during thymocyte differentiation. Dephosphorylation by protein phosphatase 1 regulates stability and pericentromeric heterochromatin location. Phosphorylated in both lymphoid and non-lymphoid tissues (By similarity). Phosphorylation at Ser-361 and Ser-364 downstream of SYK induces nuclear translocation. Sumoylated. Simultaneous sumoylation on the 2 sites results in a loss of both HDAC-dependent and HDAC-independent repression. Has no effect on pericentromeric heterochromatin location. Desumoylated by SENP1. Polyubiquitinated.
細胞内局在	Cytoplasm; Nucleus. In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events and Nucleus. In resting lymphocytes, distributed diffusely throughout the nucleus. Localizes to pericentromeric heterochromatin in proliferating cells. This localization requires DNA binding which is regulated by phosphorylation / dephosphorylation events (By similarity).
製品の状態	There are 7 isoforms produced by alternative splicing.

画像



All lanes : Anti-Ikaros antibody [EPR13790] ([ab191394](#)) at 1/10000 dilution

Lane 1 : Wild-type Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lane 2 : IKZF1 knockout Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lane 3 : Daudi (Human Burkitt's lymphoma cell line) whole cell lysate

Lane 4 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 58 kDa

Observed band size: 50-70 kDa

This data was developed using the same antibody clone in a different buffer formulation (**ab191394**).

Lanes 1 - 4: Merged signal (red and green). Green - **ab191394** observed at 50-70 kDa. Red - loading control **ab8245** (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.


ab191394 was shown to react with Ikaros in wild-type Jurkat cells in western blot with loss of signal observed in IKZF1 knockout sample. Wild-type and IKZF1 knockout Jurkat cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with **ab191394** and **ab8245** (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 10000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-Ikaros antibody
[EPR13790] - BSA and Azide free (ab206645)

Intracellular flow cytometric analysis of MOLT4 cells (paraformaldehyde-fixed, 2%) labeling Ikaros with **ab191394** at 1/130 dilution (red) or a rabbit IgG (negative) (green), followed by Goat anti rabbit IgG (FITC) secondary at 1/150 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab191394**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ikaros antibody [EPR13790] - BSA and Azide free (ab206645)

This IHC data was generated using the same anti-Ikaros antibody clone, EPR13790, in a different buffer formulation (cat# [ab191394](#)).

Immunohistochemical analysis of paraffin-embedded Human thymus tissue labeling Ikaros with [ab191394](#) at 1/1400 dilution followed by pre-diluted HRP Polymer for Rabbit IgG secondary antibody and counter-stained with Hematoxylin.
Inset: Negative control: using PBS instead of primary antibody.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ikaros antibody [EPR13790] - BSA and Azide free (ab206645)

This IHC data was generated using the same anti-Ikaros antibody clone, EPR13790, in a different buffer formulation (cat# [ab191394](#)).

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling Ikaros with [ab191394](#) at 1/1400 dilution followed by pre-diluted HRP Polymer for Rabbit IgG secondary antibody and counter-stained with Hematoxylin.
Inset: Negative control: using PBS instead of primary antibody.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Ikaros antibody [EPR13790] - BSA and Azide free (ab206645)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors