

# Anti-IKB alpha (phospho Y305) antibody ab24784

★★★★☆ [1 Abreviews](#) [2 References](#) [画像数 1](#)

### 製品の概要

製品名	Anti-IKB alpha (phospho Y305) antibody
製品の詳細	Rabbit polyclonal to IKB alpha (phospho Y305)
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide corresponding to Human IKB alpha (phospho Y305) conjugated to Keyhole Limpet Haemocyanin (KLH). Synthetic peptide conjugated to KLH corresponding to amino acid residues around tyrosine 305 of human IKB alpha. This peptide sequence has low homology to other IKB proteins.
ポジティブ・コントロール	A431 and Jurkat cells treated with pervanadate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.05% Sodium azide Constituents: PBS, 50% Glycerol, 0.1% BSA
精製度	Immunogen affinity purified
特記事項(精製)	This antibody was cross-adsorbed to phospho-tyrosine coupled to agarose then affinity purified using phospho IKB alpha (Tyr-305) peptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab24784の使用に適用されず**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★☆ (1)	1/500. Detects a band of approximately 38 kDa.

## ターゲット情報

機能	Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.
関連疾患	Ectodermal dysplasia, anhidrotic, with T-cell immunodeficiency autosomal dominant
配列類似性	Belongs to the NF-kappa-B inhibitor family. Contains 5 ANK repeats.
翻訳後修飾	Phosphorylated; disables inhibition of NF-kappa-B DNA-binding activity. Phosphorylation at positions 32 and 36 is prerequisite to recognition by UBE2D3 leading to polyubiquitination and subsequent degradation. Sumoylated; sumoylation requires the presence of the nuclear import signal. Sumoylation blocks ubiquitination and proteasome-mediated degradation of the protein thereby increasing the protein stability. Monoubiquitinated at Lys-21 and/or Lys-22 by UBE2D3. Ubiquitin chain elongation is then performed by CDC34 in cooperation with the SCF(FBXW11) E3 ligase complex, building ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. The resulting polyubiquitination leads to protein degradation. Also ubiquitinated by SCF(BTRC) following stimulus-dependent phosphorylation at Ser-32 and Ser-36. Deubiquitinated by porcine reproductive and respiratory syndrome virus Nsp2 protein, which thereby interferes with NFKBIA degradation and impairs subsequent NF-kappa-B activation.
細胞内局在	Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm by a nuclear localization signal (NLS) and a CRM1-dependent nuclear export.

## 画像



Western blot analysis of A431 cells treated with pervanadate (1 mM) for 30 minutes (30 ug/lane). Blots were probed with anti-IkappaBalpha (lane 1), anti-phospho-IkappaBalpha (Tyr-42) (**ab24783**; lanes 2-5), or anti-phospho-IkappaBalpha (Tyr-305) (ab24784; lanes 6-9). Both anti-phospho-IkappaBalpha (Tyr-42) and anti-phospho-IkappaBalpha (Tyr-305) were used in the presence of no blocking peptide (lane 2 & 6), phospho-IkappaBalpha (Tyr-42) peptide (lane 3 & 8), phospho-IkappaBalpha (Tyr-305) peptide (lane 4 & 7), or BSA conjugated to phospho-tyrosine (lane 5 & 9). Peptides and BSA-pTyr were used at 1 ug/ml.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors