

Anti-IKZF3 antibody [EPR9342(B)] ab139408

リコンビナント **RabMAb**

9 References [画像数 9](#)

製品の概要

製品名	Anti-IKZF3 antibody [EPR9342(B)]
製品の詳細	Rabbit monoclonal [EPR9342(B)] to IKZF3
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IHC-P, ICC/IF
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide within Human IKZF3 aa 1-100 (N terminal). The exact sequence is proprietary. (Peptide available as ab184024)
ポジティブ・コントロール	WB: Recombinant human IKZF3 protein (ab132618); Human thymus, Jurkat, Raji, and Ramos cell lysates. IHC-P: Human and rat spleen tissues and mouse cardiac muscle tissue. ICC/IF: Jurkat cells. Flow Cyt (intra): Ramos cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR9342(B)

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab139408の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/10 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/20000. Predicted molecular weight: 58 kDa. For unpurified use at 1/1000 - 1/10000.
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/250 - 1/500.
ICC/IF		1/100. For unpurified use at 1/250 - 1/500.

ターゲット情報

機能	Transcription factor that plays an important role in the regulation of lymphocyte differentiation. Plays an essential role in regulation of B-cell differentiation, proliferation and maturation to an effector state. Involved in regulating BCL2 expression and controlling apoptosis in T-cells in an IL2-dependent manner.
組織特異性	Expressed most strongly in peripheral blood leukocytes, the spleen, and the thymus.
配列類似性	Belongs to the Ikaros C2H2-type zinc-finger protein family. Contains 6 C2H2-type zinc fingers.
翻訳後修飾	Phosphorylation on tyrosine residues induced by IL2 is required for dissociation from HRAS and nuclear translocation of IKZF3 in T-cells. Phosphorylation on tyrosine residues induced by IL4 is required for dissociation from Bcl-X(L) in T-cells.
細胞内局在	Nucleus. Cytoplasm.

画像



All lanes : Anti-IKZF3 antibody [EPR9342(B)] (ab139408) at 1/20000 dilution (purified)

Lane 1 : Raji (Human Burkitt's lymphoma cell line) cell lysate

Lane 2 : Ramos (Human Burkitt's lymphoma cell line) cell lysate

Lane 3 : Jurkat (Human T cell leukemia cell line from peripheral blood) cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Peroxidase conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 58 kDa

Observed band size: 70 kDa

Blocking/Dilution buffer and concentration: 5% NFDM/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat spleen tissue labeling IKZF3 with purified ab139408 at 1/100.

Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500).

Negative control using PBS instead of primary antibody.

Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)




Immunocytochemistry/ Immunofluorescence - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)

Immunocytochemistry/Immunofluorescence analysis of Jurkat (Human T cell leukemia cell line from peripheral blood) cells labeling IKZF3 (green) with purified ab139408 at 1/100.

Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.


Control: Secondary antibody **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).



Flow Cytometry (Intracellular) - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)

Intracellular Flow Cytometry analysis of Ramos (Human Burkitt's lymphoma cell line) cells labeling IKZF3 with purified ab139408 at 1/90 (red).

Cells were fixed with 2% paraformaldehyde. An FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabeled control, cells without incubation with primary and secondary antibodies.



All lanes : Anti-IKZF3 antibody [EPR9342(B)] (ab139408) at 1/1000 dilution (unpurified)

Lane 1 : Human thymus cell lysate

Lane 2 : Jurkat (Human T cell leukemia cell line from peripheral blood) cell lysate

Lane 3 : Raji (Human Burkitt's lymphoma cell line) cell lysate


Lane 4 : Ramos (Human Burkitt's lymphoma cell line) cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labeled goat anti-rabbit at 1/2000 dilution

Predicted band size: 58 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human spleen tissue labeling IKZF3 with purified ab139408 at 1/100.

Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500).

Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse cardiac muscle tissue labeling IKZF3 with purified ab139408 at 1/100.

Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, an HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500).

Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IKZF3 antibody [EPR9342(B)] (ab139408)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human spleen tissue labeling IKZF3 with unpurified ab139408 at 1/250 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Anti-IKZF3 antibody [EPR9342(B)] (ab139408)

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors