# abcam

## Product datasheet

## Anti-IL-6 antibody [1.2-2B11-2G10] ab9324

★★★★★ 21 Abreviews 265 References 画像数 4

#### 製品の概要

製品名 Anti-IL-6 antibody [1.2-2B11-2G10]

製品の詳細 Mouse monoclonal [1.2-2B11-2G10] to IL-6

由来種 Mouse

アプリケーション 適用あり: IHC-P, Sandwich ELISA, WB

種交差性 交差種: Rat, Human

免疫原 Recombinant full length protein corresponding to Human IL-6 aa 29-212. (Peptide available as

ab9627).

Database link: P05231

ポジティブ・コントロール IHC-P: Human liver tumor and prostate tumor tissues; WB: Human spleen and lung tissue lysates,

Rat spleen and lung tissue lysates, Recombinant human IL6 protein (ab9627).

特記事項 Although some customers have been successful in IHC we no longer batch test in this application.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Lyophilized:Reconstitute with 500µl of sterile water. Please note that if you receive this product in

liquid form it has already been reconstituted as described and no further reconstitution is

necessary.

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

精製度 Protein A purified

**ポリ/モノ** モノクローナル

**クローン名** 1.2-2B11-2G10

アイソタイプ lgG2a

1

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab9324の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	<b>★★★★</b> <u>(12)</u>	Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
Sandwich ELISA		Use at an assay dependent concentration. In a sandwich ELISA (assuming 100µl/well), a concentration of 1.0-2.0 µg/ml of this antibody will detect at least 1000pg/ml of Recombinant Human IL-6 when used with <a href="mailto:ab271236">ab271236</a> detection antibody at a concentration of approximately 0.10-0.20 µg/ml.
WB	★★★★☆ (3)	Use at an assay dependent concentration.  To detect human IL-6 by Western blot analysis this antibody can be used at a concentration of 0.50-1.0 µg/mL. When used in conjunction with compatible secondary reagents the detection limit for recombinant human IL-6 is 1.0-2.0 ng/lane, under reducing conditions and non-reducing conditions.

#### ターゲット情報

#### 機能

Cytokine with a wide variety of biological functions. It is a potent inducer of the acute phase response. Plays an essential role in the final differentiation of B-cells into lg-secreting cells Involved in lymphocyte and monocyte differentiation. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation Acts on B-cells, T-cells, hepatocytes, hematopoeitic progenitor cells and cells of the CNS. Also acts as a myokine. It is discharged into the bloodstream after muscle contraction and acts to increase the breakdown of fats and to improve insulin resistance.

#### 関連疾患

Genetic variations in IL6 are associated with susceptibility to rheumatoid arthritis systemic juvenile (RASJ) [MIM:604302]. An inflammatory articular disorder with systemic-onset beginning before the age of 16. It represents a subgroup of juvenile arthritis associated with severe extraarticular features and occasionally fatal complications. During active phases of the disorder, patients display a typical daily spiking fever, an evanescent macular rash, lymphadenopathy, hepatosplenomegaly, serositis, myalgia and arthritis.

Note=A IL6 promoter polymorphism is associated with a lifetime risk of development of Kaposi sarcoma in HIV-infected men.


配列類似性 Belongs to the IL-6 superfamily.

N- and O-glycosylated.

細胞内局在 Secreted.


#### 画像

翻訳後修飾



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-IL-6 antibody [1.2-2B11-2G10] (ab9324)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human liver tumor tissue labelling IL-6 with ab9324 at 1  $\mu$ g/mL (45 minute incubation at room temperature). Heat mediated antigen retrieval was performed using a buffer at pH 6. An HRP-labeled polymer detection system was used with a DAB chromogen.



Western blot - Anti-IL-6 antibody [1.2-2B11-2G10] (ab9324)

**All lanes :** Anti-IL-6 antibody [1.2-2B11-2G10] (ab9324) at 1/2000 dilution

Lane 1: Human Spleen
Lane 2: Human Lung
Lane 3: Rat Spleen
Lane 4: Rat Lung

Lysates/proteins at 20 µg per lane.

## Secondary


All lanes: 800CW Goat Anti-Mouse IgG at 1/10000 dilution

Performed under reducing conditions.

Observed band size: 17,25 kDa


Additional bands at: 50 kDa (possible multimer)

This antibody was raised against an immunogen that is predicted to recognize the glycosylated form of IL6 as well as the IL6delta4 splice variant. The predicted molecular weights are 25 kDa and 17 kDa respectively. The band observed at 50 kDa may represent multimers of IL6 as reported in the literature.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-IL-6 antibody [1.2-2B11-2G10] (ab9324)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human prostate tumor tissue labelling IL-6 with ab9324 at 1  $\mu$ g/mL (45 minute incubation at room temperature). Heat mediated antigen retrieval was performed using a buffer at pH 6. An HRP-labeled polymer detection system was used with a DAB chromogen.



ab9324 detecting IL-6 by sandwich ELISA.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors