abcam

Product datasheet

Anti-iNOS antibody ab3523

★★★★ 33 Abreviews 313 References 画像数5

製品の概要

製品名 Anti-iNOS antibody

製品の詳細 Rabbit polyclonal to iNOS

由来種 Rabbit

特異性 This antibody detects iNOS. It does not detect other NOS isoforms. By western blot, this antibody

> detects an ~135 kDa protein representing recombinant human iNOS. By western blot, this antibody also detects purified recombinant mouse iNOS, mouse iNOS from cytokine stimulated

RAW 264.7 cells.

アプリケーション 適用あり: IHC-P, WB, ICC/IF

種交差性 交差種: Mouse, Human

免疫原 Synthetic peptide corresponding to Mouse iNOS aa 1-100.

Database link: P29477

Run BLAST with Run BLAST with

ポジティブ・コントロール

ICC: A549, NIH/3T3 cells. IHC-P: Human heart tissue.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

> Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー Preservative: 0.05% Sodium azide

Constituents: 0.1% BSA, 99% PBS

精製度 Immunogen affinity purified

ポリモノ ポリクローナル

アイソタイプ ΙgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab3523の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★ ☆ <u>(12)</u>	1/20.
WB	★★★★★ (15)	1/200 - 1/1000. Predicted molecular weight: 131 kDa.
ICC/IF	★★★★★ (2)	1/50.

ターゲット情報

機能
Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such COX2.


組織特異性
Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in

the platelets.

配列類似性 Belongs to the NOS family.

Contains 1 FAD-binding FR-type domain. Contains 1 flavodoxin-like domain.

画像



All lanes : Anti-iNOS antibody (ab3523) at 1/1000 dilution

Lane 1: RAW264 whole cell lysate untreated

Lane 2: RAW264 whole cell lysate untreated stimulated with LPS

at 1 µg/mL for 16 hours

Lysates/proteins at 20 µg per lane.

Secondary

Lane 1 : Goat anti-Rabbit lgG (Heavy Chain) Superclonal Secondary Antibody, HRP conjugate at 1/1000 dilution

 $\textbf{Lane 2:} \ \ \textbf{Goat anti-Rabbit IgG (Heavy Chain) HRP conjugate at}$

1/1000 dilution

Predicted band size: 131 kDa

Using 4-20% Tris-Glycine polyacrylamide gel and transferred to a nitrocellulose membrane, blocked with 5% Milk in TBST for at least 1 hour. The membrane was probed with ab3523 at 4°C overnight on a rocking platform, washed in TBST, and probed with the secondary antibody for 1 hour.

All lanes: Anti-iNOS antibody (ab3523) at 1/1000 dilution

Lane 1: Whole cell lysate of RAW 264.7

Lane 2: Whole cell lysate of RAW 264.7 treated with LPS

Lysates/proteins at 30 µg per lane.



All lanes : Goat anti-Rabbit IgG (Heavy Chain) Superclonal™ Recombinant Secondary Antibody, HRP at 1/4000 dilution


Predicted band size: 131 kDa

This was run using 4-12% Bis-Tris Protein Gel and a Nitrocellulose membrane.


Observed band at 130kDa in LPS treated RAW 264.7 cells, and an uncharacterized band at ~60kDa.

Immunohistochemistry analysis of human heart tissue stained for iNOS without (negative control) or using ab3523 at 1/200 dilution overnight at 4°C in a humidified chamber, followed biotin-conjugated secondary antibody and SA-HRP, followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor.


To expose target proteins, heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature.








Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-iNOS antibody (ab3523)



Immunocytochemistry/ Immunofluorescence - AntiiNOS antibody (ab3523)

Immunofluorescence analysis of A549 (human lung carcinoma cell line) whole cells labelling iNOS (Left panel: green) without (control) or using ab3523 at 1/20 dilution overnight at 4°C, followed DyLight-488 conjugated secondary antibody. Counter stain: F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Images were taken at 60X magnification.

Cells were grown on chamber slides and fixed with formaldehyde prior to staining.



Immunocytochemistry/ Immunofluorescence - AntiiNOS antibody (ab3523)

Immunofluorescence analysis of NIH/3T3 (mouse embryo fibroblast cell line) whole cells labelling iNOS (Left panel: green) without (control) or using ab3523 at 1/20 dilution overnight at 4°C, followed DyLight-488 conjugated secondary antibody. Counter stain: F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Images were taken at 60X magnification.

Cells were grown on chamber slides and fixed with formaldehyde prior to staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors