

Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] ab269948

リコンビナント **RabMAb**

1 References [画像数 7](#)

製品の概要

製品名	Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12]
製品の詳細	Rabbit monoclonal [EPR23244-12] to Junctional Adhesion Molecule 1/JAM-A
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB, Flow Cyt, IP 適用なし: IHC-P
種交差性	交差種: Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human lung tissue lysate. HT-29, HUVEC and TF-1 whole cell lysate. ICC/IF: HT-29 cells. Flow Cyt: HT-29 cells, human B lymphocytes and human monocytes. IP: HT-29 whole cell lysate. Human lung tissue lysate.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified

ポリ/モノ	モノクローナル
クローン名	EPR23244-12
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab269948の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
ICC/IF		1/50.
WB		1/1000. Predicted molecular weight: 32 kDa.
Flow Cyt		1/500.
IP		1/30.

追加情報 Is unsuitable for IHC-P.

ターゲット情報

機能	Seems to plays a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3. The association of the PARD6-PARD3 complex may prevent the interaction of PARD3 with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier. Involved in platelet activation. In case of orthoreovirus infection, serves as receptor for the virus.
配列類似性	Belongs to the immunoglobulin superfamily. Contains 2 Ig-like V-type (immunoglobulin-like) domains.
翻訳後修飾	N-glycosylated.
細胞内局在	Cell junction > tight junction. Cell membrane. Localized at tight junctions of both epithelial and endothelial cells.

画像



All lanes : Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948) at 1/1000 dilution

Lane 1 : Human lung lysate

Lane 2 : HT-29 (human colorectal adenocarcinoma epithelial cell) whole cell lysate

Lane 3 : THP-1 (human monocytic leukemia monocyte) whole cell lysate

Lane 4 : HUVEC (human umbilical vein endothelial cell) whole cell lysate

Lane 5 : MOLT-4 (human lymphoblastic leukemia t lymphoblast) whole cell lysate

Lane 6 : TF-1 (human erythroleukemia erythroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/50000 dilution

Predicted band size: 32 kDa


Observed band size: 35 kDa

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID:10698320, 10753840).

Low expression control: THP-1 and Molt-4 (PMID:10698320).

Exposure time: Lanes 1-5: 15 seconds; Lane 6:70 seconds.



Immunoprecipitation - Anti-Junctional Adhesion
Molecule 1/JAM-A antibody [EPR23244-12]
(ab269948)

Junctional Adhesion Molecule 1/JAM-A was immunoprecipitated from 0.35 mg human lung lysate 10µg with ab269948 at 1/30 dilution (2µg in 0.35 mg lysates). Western blot was performed on the immunoprecipitate using ab269948 at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.


Lane 1: Human lung lysate 10µg.

Lane 2: ab269948 IP in human lung lysate.

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab269948 in human lung lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds.



Immunoprecipitation - Anti-Junctional Adhesion
Molecule 1/JAM-A antibody [EPR23244-12]
(ab269948)

Junctional Adhesion Molecule 1/JAM-A was immunoprecipitated from 0.35 mg HT-29 (Human colorectal adenocarcinoma epithelial cell) whole cell lysate 10µg with ab269948 at 1/30 dilution (2µg in 0.35 mg lysates). Western blot was performed on the immunoprecipitate using ab269948 at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.


Lane 1: HT-29 whole cell lysate 10µg.

Lane 2: ab269948 IP in HT-29 whole cell lysate.

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab269948 in HT-29 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds.




Flow Cytometry - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Flow cytometric analysis of human B lymphocytes (Left) / human monocytes (Right) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/500 compared with a Rabbit monoclonal IgG ([ab172730](#)) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) at 1/2000 was used as the secondary antibody.

Human peripheral blood mononuclear cell (PBMC) co-stained with anti-CD19 conjugated to PE-Cy7 and anti-CD14 conjugated to BV510. JAM-A expression on B lymphocytes (CD19+) and monocytes (CD14+) population are shown respectively.


Gated on viable cells.



Flow Cytometry - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Flow cytometric analysis of HT-29 (Human colorectal adenocarcinoma epithelial cell) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/500 compared with a Rabbit monoclonal IgG ([ab172730](#)) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) at 1/2000 was used as the secondary antibody.

Gated on viable cells.



Immunocytochemistry/ Immunofluorescence - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HT-29 (human colorectal adenocarcinoma epithelial cell) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/50 dilution, followed by [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing membranous staining in HT-29 cell line is observed. [ab195889](#) Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: [ab150077](#) AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Junctional Adhesion Molecule 1/JAM-A
antibody [EPR23244-12] (ab269948)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors