

Anti-Ku70 antibody [EPR4027] ab92450

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **14 References** 画像数 10

製品の概要

製品名	Anti-Ku70 antibody [EPR4027]
製品の詳細	Rabbit monoclonal [EPR4027] to Ku70
由来種	Rabbit
アプリケーション	適用あり: WB, IP, IHC-P, ICC/IF, Flow Cyt (Intra)
種交差性	交差種: Human
免疫原	Synthetic peptide within Human Ku70 aa 500-600. The exact sequence is proprietary. Database link: P12956
ポジティブ・コントロール	WB: A549, 293T, A431, and HeLa lysates. IHC-P: human colon carcinoma, tonsil and testis tissue. ICC/IF: HeLa cells. Flow Cyt (intra): HeLa cells. IP: HeLa cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified

ポリ/モノ	モノクローナル
クローン名	EPR4027
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab92450の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/1000 - 1/10000. Predicted molecular weight: 70 kDa.
IP		1/20.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Antigen retrieval and the use of an HRP/AP polymerized secondary antibody is recommended for enhanced staining. See <u>IHC antigen retrieval protocols</u> .
ICC/IF		1/100 - 1/250.
Flow Cyt (Intra)		1/10 - 1/100. <u>ab172730</u> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

ターゲット情報


機能	Single stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. Required for osteocalcin gene expression. Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks. 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription.
配列類似性	Belongs to the ku70 family. Contains 1 Ku domain. Contains 1 SAP domain.
発生段階	Expression does not increase during promyelocyte differentiation.
翻訳後修飾	Phosphorylation by PRKDC may enhance helicase activity. Phosphorylation of Ser-51 does not

affect DNA repair.

細胞内局在

Nucleus. Chromosome.

画像



All lanes : Anti-Ku70 antibody [EPR4027] (ab92450) at 1/5000 dilution (Purified)

Lane 1 : A549 (Human lung carcinoma epithelial cell) whole cell lysates

Lane 2 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates


Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 70 kDa

Observed band size: 70 kDa




Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Ku70 with purified ab92450 at 1/250 dilution (0.46 µg/ml). Cells were fixed in 100% Methanol and permeabilized with None. Cells were counterstained with **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ku70 antibody [EPR4027] (ab92450)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon carcinoma tissue sections labeling Ku70 with purified ab92450 at 1/250 dilution (0.46 µg/ml). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 1 (pH 6.0). Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody.

Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-Ku70 antibody [EPR4027] (ab92450)

All lanes : Anti-Ku70 antibody [EPR4027] (ab92450) at 1/1000 dilution ((unpurified))

Lane 1 : HeLa cell lysate

Lane 2 : 293T cell lysate

Lane 3 : A549 cell lysate


Lane 4 : A431 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 70 kDa



Flow Cytometry (Intracellular) - Anti-Ku70 antibody
[EPR4027] (ab92450)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Ku70 with purified ab92450 at 1/20 dilution (5µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunoprecipitation - Anti-Ku70 antibody
[EPR4027] (ab92450)

ab92450 (purified) at 1/20 dilution (0.5ug) immunoprecipitating Ku70 in HeLa whole cell lysate.


Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10ug

Lane 2 (+): ab92450 & HeLa whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab92450 in HeLa whole cell lysate


For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.




Immunocytochemistry/ Immunofluorescence - Anti-Ku70 antibody [EPR4027] (ab92450)

Immunofluorescence staining of HeLa cells using unpurified ab92450 at 1/100 dilution.



Flow Cytometry (Intracellular) - Anti-Ku70 antibody [EPR4027] (ab92450)

Overlay histogram showing HeLa cells stained with unpurified ab92450 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab92450, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HeLa cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ku70 antibody [EPR4027] (ab92450)

Paraffin embedded Human testis tissue (A) or Human tonsil tissue (B) were labelled with unpurified ab92450 at 1/100 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Ku70 antibody [EPR4027] (ab92450)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors