


### Anti-Lamin B1 antibody - Nuclear Envelope Marker ab16048

KO 評価済

★★★★★ [107 Abreviews](#) [1046 References](#) [画像数 12](#)

#### 製品の概要

製品名	Anti-Lamin B1 antibody - Nuclear Envelope Marker
製品の詳細	Rabbit polyclonal to Lamin B1 - Nuclear Envelope Marker
由来種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, WB, IHC-P
種交差性	<b>交差種:</b> Mouse, Rat, Human <b>交差が予測される動物種:</b> Chicken, Pig, Xenopus laevis, Indian muntjac, Zebrafish 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as <a href="#">ab16375</a> )
ポジティブ・コントロール	ICC/IF: HepG2 cells WB: HeLa, PC12 and NIH/3T3 whole cell lysate. Wild type HAP1 whole cell lysate. Wild type HAP1 nuclear lysate. Human Pancreatic cell line whole cell lysate. IHC-P: Human liver tissue. Human infantile fibromatosis tissue.
特記事項	<p>Lamin B1 and Lamin B antibodies are extremely useful as nuclear loading controls for use with nuclear extracts. When using Lamin B1 antibodies as nuclear loading controls, be aware that in apoptotic cells Lamin B1 is cleaved (Kottke TJ et al.). Lamin B1 will also be removed from a nuclear prep if the nuclear membranes are spun out. This antibody was designed to be a nuclear loading control however it has not yet been tested in appropriate lysates.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40

Preservative: 0.02% Sodium azide

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

#### 精製度

Immunogen affinity purified

#### 一次抗体 備考

Lamin B1 and Lamin B antibodies are extremely useful as nuclear loading controls for use with nuclear extracts. When using Lamin B1 antibodies as nuclear loading controls, be aware that in apoptotic cells Lamin B1 is cleaved (Kottke TJ et al.). Lamin B1 will also be removed from a nuclear prep if the nuclear membranes are spun out. This antibody was designed to be a nuclear loading control however it has not yet been tested in appropriate lysates.

#### ポリ/モノ

ポリクローナル

#### アイソタイプ

IgG

#### アプリケーション

##### The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab16048の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (27)	Use a concentration of 0.1 µg/ml.
WB	★★★★★ (66)	Use a concentration of 0.1 µg/ml. Detects a band of approximately 68 kDa (predicted molecular weight: 66 kDa). We recommend <b>Goat anti-Rabbit IgG H&amp;L (IRDye® 800CW) preadsorbed (ab216773)</b> .
IHC-P	★★★★★ (5)	Use a concentration of 0.1 - 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

#### ターゲット情報

##### 機能

Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin.

##### 関連疾患

Defects in LMNB1 are the cause of leukodystrophy demyelinating autosomal dominant adult-onset (ADLD) [MIM:169500]. ADLD is a slowly progressive and fatal demyelinating leukodystrophy, presenting in the fourth or fifth decade of life. Clinically characterized by early autonomic abnormalities, pyramidal and cerebellar dysfunction, and symmetric demyelination of the CNS. It differs from multiple sclerosis and other demyelinating disorders in that neuropathology shows preservation of oligodendroglia in the presence of subtotal demyelination and lack of astrogliosis.

##### 配列類似性



Belongs to the intermediate filament family.

##### 翻訳後修飾

B-type lamins undergo a series of modifications, such as farnesylation and phosphorylation. Increased phosphorylation of the lamins occurs before envelope disintegration and probably plays

	a role in regulating lamin associations.
細胞内局在	Nucleus inner membrane.

画像



Immunocytochemistry analysis of formaldehyde-fixed NP40-permeabilized Mouse vascular smooth muscle cells staining with ab16048 at 1/100. Secondary antibody was Alexa Fluor® 488 donkey Anti-rabbit at 1/500 dilution. Samples were incubated with the primary antibody with 3% BSA in PBS for 12 hours at 4°C. Blocking was done using 3% BSA for 1 hour at 21°C.

**All lanes** : Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048) at 1 µg/ml

**Lane 1** : HeLa whole cell lysate

**Lane 2** : PC12 whole cell lysate

**Lane 3** : NIH 3T3 whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

**All lanes** : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/50000 dilution

**Predicted band size:** 66 kDa


Blocking buffer: 3% Milk

Gel type: MOPS

Exposure Time: 2 minutes


Observed band size: 73 kDa

Additional bands: 46 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

Immunohistochemical analysis of formalin-fixed paraffin-embedded human liver labelling lamin B1 with ab16048 at a concentration of 0.2µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5). ab16048 anti lamin B1 antibody was incubated at 37°C for 16min. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.



Western blot - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

**Lane 1:** Wild type HAP1 whole cell lysate (20 µg)


**Lane 2:** empty lane

**Lane 3:** KO HAP1 LMNB1 whole cell lysate (20 µg)

**Lane 4:** empty lane

**Lanes 1 - 4:** Merged signal (red and green). Green - ab16048 observed at 70 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab16048 was shown to specifically react with LMNB1 (Lamin B1) in wild type HAP1 cells. No band was observed when LMNB1 (Lamin B1) knockout samples were used. Ab16048 LMNB1 (Lamin B1) and **ab8245** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 0.1 µg per mL and 1/10000 dilution respectively. Blots were developed with **Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773)** and **Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776)** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

ab16048 staining Lamin B1 in HepG2 cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab16048 at 0.1 µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Also suitable in cells fixed with 100% methanol (5 min).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

IHC image of Lamin B1 staining in Human normal Liver formalin fixed paraffin embedded tissue section\*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab16048, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Western blot - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

**All lanes :** Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048) at 1 µg/ml

**Lane 1 :** Wild-type HAP1 nuclear lysate

**Lane 2 :** Lamin B1 knockout HAP1 nuclear lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) at 1/10000 dilution

Performed under reducing conditions.

**Predicted band size:** 66 kDa

**Observed band size:** 68 kDa


**Lane 1:** Wild-type HAP1 nuclear lysate (10 µg)

**Lane 2:** Lamin B1 knockout HAP1 nuclear lysate (10 µg)

**Lanes 1 and 2:** Green signal from target - ab16048 observed at 68 kDa. Red signal from loading control - **ab10799** observed at 18 kDa.

ab16048 was shown to specifically react with lamin B1 in wild-type HAP1 cells. No band was observed knockout samples were used. Wild-type and lamin B1 knockout samples were subjected to SDS-PAGE. ab16048 and **ab10799** (loading control to histone H3 at 0.1µg/mL) were both incubated overnight at 4°C. Blots were developed with **Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773)** and **Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776)** secondary antibodies at 1/10,000 dilution for 1 h at room temperature before imaging.





Western blot - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

**All lanes :** Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048) at 1/1000 dilution

**Lane 1 :** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

**Lane 2 :** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate with Mouse Lamin B1 peptide ([ab16375](#)) at 1 µg/ml

Lysates/proteins at 20 µg per lane.


#### Secondary

**All lanes :** Alexa fluor goat polyclonal to Rabbit IgG at 1/10000 dilution

Performed under reducing conditions.

**Predicted band size:** 66 kDa

**Observed band size:** 68-70 kDa



Western blot - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

This image is courtesy of an anonymous Abreview

Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048) at 1/1000 dilution + Human Pancreatic cell line - whole cell lysate at 20 µg

#### Secondary

HRP conjugated goat anti-rabbit antibody at 1/2000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 66 kDa

**Observed band size:** 68 kDa


**Exposure time:** 30 seconds



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

This image is courtesy of an anonymous Abreview

ab16048 staining Lamin B1 in human infantile fibromatosis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 1% FBS/BSA for 3 hours at room temperature; antigen retrieval was by heat mediation in Tris pH9. Samples were incubated with primary antibody (1/100 in TBS + 1% BSA + 1% FBS) for 16 hours. An undiluted HRP-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

Image courtesy of Marilena Ciciarello and Patrizia Lavia, University 'La Sapienza' CNR, Italy

Human and mouse cells stained with ab16048 (1/500). The cells were fixed and permeabilized in 4% formaldehyde, 0.2% Tritonm X100 for 10 minutes at room temperature, then washed 3x in PBS.


A: HeLa cells + ab16048 (green)

B: HeLa cells counterstained with DAPI (blue)

C: 3T3 cells + ab16048 (green)

D: 3T3 cells counterstained with DAPI (blue)





Human and mouse cells stained with ab16048 (1/500). The cells were fixed in 100% methanol for 6 minutes at -20°C, then washed once in PBS.

A: HeLa cells + ab16048 (green)

B: HeLa cells counterstained with DAPI (blue)

C: 3T3 cells + ab16048 (green)

D: 3T3 cells counterstained with DAPI (blue)

Immunocytochemistry/ Immunofluorescence - Anti-Lamin B1 antibody - Nuclear Envelope Marker (ab16048)

Image courtesy of Marilena Ciciarello and Patrizia Lavia, University 'La Sapienza' CNR, Italy

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors