

Anti-LAMP2 antibody [EPR19531] ab199947

KO 評価済 リコンビナント RabMAb

2 References 画像数 7

製品の概要

製品名	Anti-LAMP2 antibody [EPR19531]
製品の詳細	Rabbit monoclonal [EPR19531] to LAMP2
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human fetal kidney, fetal spleen and placenta lysates; HeLa, THP-1, HepG2, HEK-293, JAR and Jurkat whole cell lysates. IHC-P: Human liver and breast cancer tissues.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR19531
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab199947の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB		1/1000. Detects a band of approximately 110-130 kDa (predicted molecular weight: 45 kDa).

ターゲット情報

機能	Implicated in tumor cell metastasis. May function in protection of the lysosomal membrane from autodigestion, maintenance of the acidic environment of the lysosome, adhesion when expressed on the cell surface (plasma membrane), and inter-and intracellular signal transduction. Protects cells from the toxic effects of methylating mutagens.
組織特異性	Isoform LAMP-2A is highly expressed in placenta, lung and liver, less in kidney and pancreas, low in brain and skeletal muscle. Isoform LAMP-2B is highly expressed in skeletal muscle, less in brain, placenta, lung, kidney and pancreas, very low in liver.
関連疾患	Defects in LAMP2 are the cause of Danon disease (DAND) [MIM:300257]; also known as glycogen storage disease type 2B (GSD2B). DAND is a lysosomal glycogen storage disease characterized by the clinical triad of cardiomyopathy, vacuolar myopathy and mental retardation. It is often associated with an accumulation of glycogen in muscle and lysosomes.
配列類似性	Belongs to the LAMP family.
翻訳後修飾	O- and N-glycosylated; some of the 16 N-linked glycans are polylactosaminoglycans.
細胞内局在	Cell membrane. Endosome membrane. Lysosome membrane. This protein shuttles between lysosomes, endosomes, and the plasma membrane.
製品の状態	Alternative splicing produces 3 isoforms.

画像



Western blot - Anti-LAMP2 antibody [EPR19531] (ab199947)

All lanes : Anti-LAMP2 antibody [EPR19531] (ab199947) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell

Lane 2 : LAMP2 CRISPR-Cas9 edited HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 45 kDa

Observed band size: 100 kDa

False colour image of Western blot: Anti-LAMP2 antibody [EPR19531] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (**ab7291**) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab199947 was shown to bind specifically to LAMP2. A band was observed at 100 kDa in wild-type HeLa cell lysates with no signal observed at this size in LAMP2 CRISPR-Cas9 edited cell line **ab255402** (CRISPR-Cas9 edited cell lysate **ab263861**). The band observed in the CRISPR-Cas9 edited lysate lane below 100 kDa is likely to represent a truncated form of LAMP2. This has not been investigated further and the functional properties of the gene product have not been determined. To generate this image, wild-type and LAMP2 CRISPR-Cas9 edited HeLa cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were

washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



All lanes : Anti-LAMP2 antibody [EPR19531] (ab199947) at 1/1000 dilution

Lane 1 : Wild-type HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate

Lane 2 : LAMP2 knockout HEK-293 (Human epithelial cell line from embryonic kidney) whole cell lysate

Lane 3 : Hep G2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 4 : THP-1 (Human monocytic leukemia cell line) whole cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 45 kDa

Lanes 1 -4: Merged signal (red and green). Green - ab199947 observed at 45 kDa. Red - loading control, **ab130007**, observed at 130 kDa.

ab199947 was shown to specifically react with LAMP2 in wild-type HEK-293 cells as signal was lost in LAMP2 knockout cells. Wild-type and LAMP2 knockout samples were subjected to SDS-PAGE. Ab199947 and **ab130007** (Mouse anti-Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-LAMP2 antibody [EPR19531] (ab199947) at 1/1000 dilution

Lane 1 : Human fetal kidney lysate

Lane 2 : Human fetal spleen lysate

Lane 3 : Human placenta lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/100000 dilution


Predicted band size: 45 kDa

Observed band size: 110-130 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFD/MTBST.

The molecular weight observed is consistent with what has been described in the literature (PMID: 19828315) and lots of other products.



All lanes : Anti-LAMP2 antibody [EPR19531] (ab199947) at 1/1000 dilution

Lane 1 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lane 2 : THP-1 (Human monocytic leukemia cell line) whole cell lysate

Lane 3 : JAR (Human placenta choriocarcinoma cell line) whole cell lysate

Lane 4 : Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 45 kDa

Observed band size: 110-130 kDa

Exposure time: 3 minutes

Blocking/Dilution buffer: 5% NFDm/TBST.

The molecular weight observed is consistent with what has been described in the literature (PMID: 19828315) and lots of other products.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LAMP2 antibody [EPR19531] (ab199947)


Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling LAMP2 with ab199947 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Cytoplasm staining on Human liver is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LAMP2 antibody [EPR19531] (ab199947)

Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue labeling LAMP2 with ab199947 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.





Cytoplasm staining on Human breast cancer is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-LAMP2 antibody [EPR19531] (ab199947)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors