

Anti-LC3B antibody [EPR18709] - BSA and Azide free ab221794

KO 評価済 リコンビナント RabMAb

[1 Abreviews](#) [5 References](#) [画像数 12](#)

製品の概要

製品名	Anti-LC3B antibody [EPR18709] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR18709] to LC3B - BSA and Azide free
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF 適用なし: IP
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: BMDM, U-87 MG, C6 and RAW 264.7 whole cell lysates; Human brain, mouse heart, rat heart, mouse brain and rat brain lysates. ICC/IF: HeLa cells (+/- chloroquine), HAP1 cells (+/- chloroquine) (HAP1-MAP1LC3B knockout cells used as negative cell line); H9C2 rat cardiomyocytes. IHC-P: Human Cerebral Cortex tissue sections
特記事項	<p>ab221794 is the carrier-free version of ab192890.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR18709
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab221794の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Detects a band of approximately 14, 16 kDa (predicted molecular weight: 15 kDa).
IHC-P		Use a concentration of 0.1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.

追加情報 Is unsuitable for IP.

ターゲット情報

機能	Probably involved in formation of autophagosomal vacuoles (autophagosomes).
組織特異性	Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.
配列類似性	Belongs to the MAP1 LC3 family.
翻訳後修飾	The precursor molecule is cleaved by APG4B/ATG4B to form LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form LC3-II.
細胞内局在	Cytoplasm > cytoskeleton. Endomembrane system. Cytoplasmic vesicle > autophagosome membrane. LC3-II binds to the autophagic membranes.

画像



All lanes : Anti-LC3B antibody [EPR18709] - Autophagosome

Marker ([ab192890](#)) at 1/2000 dilution

Lane 1 : Wild-type HepG2 untreated control cell lysate

Lane 2 : Wild-type HepG2 Treated Chloroquine (50 uM, 16 h) cell lysate

Lane 3 : MAP1LC3B knockout HepG2 untreated control cell lysate

Lane 4 : MAP1LC3B knockout HepG2 Treated Chloroquine (50 uM, 16 h) cell lysate

Lane 5 : U-87 MG cell lysate

Lane 6 : PC-12 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 15 kDa

Observed band size: 14,16 kDa


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab192890](#)).

False colour image of Western blot: Anti-LC3B antibody [EPR18709] - Autophagosome Marker staining at 1/2000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] ([ab7291](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, [ab192890](#) was shown to bind specifically to LC3B. A band was observed at 16/14 kDa (yellow arrows) in treated wild-type HepG2 cell lysates with no signal observed at this size in MAP1LC3B knockout cell line [ab277828](#) (knockout cell lysate [ab283796](#)). To generate this image, wild-type and MAP1LC3B knockout HepG2 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Western blot - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

This data was developed using **ab192890**, the same antibody clone in a different buffer formulation. Different batches of **ab192890** were tested on U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) lysate at 0.9 µg/ml. 15 µg of lysate was loaded in each lane. Bands observed at 14, 16 kDa.




Immunocytochemistry/ Immunofluorescence - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

ab192890 staining LC3B in HeLa cells +/- Chloroquine (50 µM, 24 hours). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with **ab192890** at 1 µg/ml and **ab195889**, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 1/250 dilution (shown in pseudocolor red) followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab192890**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab192890**).

ab192890 staining LC3B in paraffin embedded human astrocytoma tissue by Immunohistochemistry. Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins. Samples were incubated with primary antibody at 1/1000 dilution for 30 mins at room temperature. Ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used as the secondary antibody. Hematoxylin was used as a counterstain. The immunostaining was performed on a Leica Biosystems BOND® RX instrument.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab192890**).

Immunohistochemical analysis of formalin fixed paraffin embedded human cortex labelling LC3B with **ab192890** at a dilution of 0.1 µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5) . **ab192890** anti LC3B antibody was incubated at 37°C for 16min. Sections were counterstained is with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.

Tissue Microarray (TMA) data for ab192890					
Normal tissue samples			Malignant tissue samples		
Human cardiac muscle	x	Human placenta	x	Human glioma	✓
Human cerebrum	✓	Human skeletal muscle	x	Human hepatocellular carcinoma	x
Human colon	x	Human skin	x	Human lung carcinoma	x
Human endometrium	x	Human spleen	x	Human ovarian carcinoma	x
Human kidney	x	Human stomach	x	Human pancreatic carcinoma	x
Human liver	x	Human testis	x	Human prostatic hyperplasia	x
Human lung	x	Human thyroid	x	Human thyroid carcinoma	x
Human mammary gland	x	Human tonsil	x		
Human pancreas	x				
		Clear cell carcinoma of human kidney	x		
		Human bladder cancer	x		
		Human breast carcinoma	x		
		Human cervical carcinoma	x		
		Human colon carcinoma	x		
		Human endometrial carcinoma	x		
		Human gastric adenocarcinoma	x		

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab192890**).

Tissue Microarrays stained for Anti-LC3B antibody [EPR18709] - Autophagosome Marker using **ab192890** in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negative (cross mark) staining per sample type tested.


Heat mediated antigen retrieval was performed with Citrate buffer (pH 6.0, Epitope Retrieval Solution 1) for 20 mins.

The section was incubated with **ab192890** for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

IHC image of LC3B staining in a section of formalin-fixed paraffin-embedded normal human cerebral cortex* performed on a Leica BOND™ system using the standard protocol **F**. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab221794, 0.1ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.




Immunocytochemistry/ Immunofluorescence - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

Clone EPR18709 (ab221794) has been successfully conjugated by Abcam. This image was generated using Anti-LC3B antibody [EPR18709] - Autophagosome Marker (Alexa Fluor® 647). Please refer to [ab225383](#) for protocol details.

[ab225383](#) staining LC3B in wild-type HAP1 cells and knockout cells, untreated and chloroquine-treated (50µM, 24 hours). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with [ab225383](#) at 0.5µg/ml (shown in red) and [ab195887](#), Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green) overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

Clone EPR18709 (ab221794) has been successfully conjugated by Abcam. This image was generated using Anti-LC3B antibody [EPR18709] - Autophagosome Marker (Alexa Fluor® 488). Please refer to [ab225382](#) for protocol details.

[ab225382](#) staining LC3B in HeLa chloroquine-treated (50µM, 24 hours) cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with [ab225453](#) at 1/100 dilution (shown in green) and [ab195889](#), Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

This image is courtesy of an Abreview submitted by Kirk McManus, Univ. of Manitoba/Cancer Care MCB.

Immunocytochemistry/ Immunofluorescence analysis of HeLa cells labeling LC3B with [ab192890](#) at 1/500 dilution. Cells were fixed in Methanol. Staining with [ab192890](#) at 1/500 was carried out for 1 hour at 22°C in PBS buffer. [ab150081](#), a Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed secondary antibody was used at 1/200 dilution. DAPI was used to counterstain.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab192890](#)).



This ICC data was generated using the same anti-LC3B antibody clone, EPR18709, in a different buffer formulation (cat# **ab192890**).

ab192890 staining LC3B in HAP1 cells (wildtype and MAP1LC3B knockout) +/- Chloroquine (50µM, 24 hours). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with **ab192890** at 1µg/ml and **ab195889**, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 1/250 dilution (shown in pseudocolor red) followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-LC3B antibody [EPR18709] - BSA and Azide free (ab221794)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors