


Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free ab183216

KO 評価済 リコンビナント RabMAb

4 References 画像数 4

製品の概要

製品名	Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free
製品の詳細	Rabbit monoclonal [MJFF3 (c69-6)] to LRRK2 - BSA and Azide free
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Human brain hippocampus tissue WB: Jurkat cells transfected with LRRK2
特記事項	<p>ab183216 is the carrier-free version of ab133475.</p> <p>Well-characterized antibodies to efficiently detect and purify LRRK2 protein are a critical need in the Parkinson's Disease (PD) research community. To help accelerate LRRK2 research, The Michael J. Fox Foundation (MJFF), working with Epitomics, Inc., has generated unique and high quality LRRK2 rabbit monoclonal antibodies to be widely available for PD research community.</p> <p>LRRK2 (Leucine-rich repeat kinase 2, dardarin) is a protein kinase belonging to the ROCO family, which is defined by the presence of a ROC (Ras/GTPase of complex proteins) domain and COR (C-terminal of Roc) region. LRRK2 exhibits kinase activity whereby it can undergo autophosphorylation and can phosphorylate generic substrates. In addition, the GTPase domain of LRRK2 can mediate GDP (guanosine-5'-diphosphate)/GTP (guanosine-5'-triphosphate) binding as well as GTP hydrolysis.</p> <p>LRRK2 is mutated in a significant number of Parkinson's disease (PD) patients. Mutations in this gene account for 4% of PD, and are observed in 1% of sporadic PD patients. Clinical symptoms of patients carrying PD-associated mutations of LRRK2 are indistinguishable from typical sporadic PD. The spectra of neuropathological features of PARK8 (type 8), the type corresponding to LRRK2, is broad and appears to encompass those associated with other familial PD cases such as PARK1 (alpha-synuclein) and PARK2 (Parkin). Patients with this gene mutation have typical relatively late onset Parkinsonism with features comparable with idiopathic PD; symptoms include asymmetric rest tremor, bradykinesia, rigidity, and a good response to 3,4-dihydroxy-L-phenylalanine (L-DOPA). The pathology of cases with LRRK2 mutations is pleomorphic.</p> <p>For more characterization data and protocols using this LRRK2 Antibody, please refer to Davies,</p>

et al. 2013. Biochemical J 453(1):101-113 [PMID: 23560750]

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

This antibody was developed with support from The Michael J. Fox Foundation.



製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	MJFF3 (c69-6)
アイソタイプ	IgG


アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab183216の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Predicted molecular weight: 286 kDa. Please check the parent abID, ab133475 , for a recommended dilution.

ターゲット情報

機能	Positively regulates autophagy through a calcium-dependent activation of the CaMKK/AMPK signaling pathway. The process involves activation of nicotinic acid adenine dinucleotide phosphate (NAADP) receptors, increase in lysosomal pH, and calcium release from lysosomes. Together with RAB29, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner. Regulates neuronal process morphology in the intact central nervous system (CNS). Plays a role in synaptic vesicle trafficking. Phosphorylates PRDX3. Has GTPase activity. May play a role in the phosphorylation of proteins central to Parkinson disease.
組織特異性	Expressed in the brain. Expressed in pyramidal neurons in all cortical laminae of the visual cortex, in neurons of the substantia nigra pars compacta and caudate putamen (at protein level). Expressed throughout the adult brain, but at a lower level than in heart and liver. Also expressed in placenta, lung, skeletal muscle, kidney and pancreas. In the brain, expressed in the cerebellum, cerebral cortex, medulla, spinal cord occipital pole, frontal lobe, temporal lobe and putamen. Expression is particularly high in brain dopaminoceptive areas.
関連疾患	Parkinson disease 8
配列類似性	Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Contains 12 LRR (leucine-rich) repeats. Contains 1 protein kinase domain. Contains 1 Roc domain. Contains 7 WD repeats.
ドメイン	The seven-bladed WD repeat region is critical for synaptic vesicle trafficking and mediates interaction with multiple vesicle-associated presynaptic proteins. The Roc domain mediates homodimerization and regulates kinase activity.
翻訳後修飾	Autophosphorylated.
細胞内局在	Membrane. Cytoplasm. Perikaryon. Mitochondrion. Golgi apparatus. Cell projection, axon. Cell projection, dendrite. Endoplasmic reticulum. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Endosome. Lysosome. Mitochondrion outer membrane. Mitochondrion inner membrane. Mitochondrion matrix. Predominantly associated with intracytoplasmic vesicular and membranous structures (By similarity). Localized in the cytoplasm and associated with cellular membrane structures. Predominantly associated with the mitochondrial outer membrane of the mitochondria. Colocalized with RAB29 along tubular structures emerging from Golgi apparatus. Localizes in intracytoplasmic punctate structures of neuronal perikarya and dendritic and axonal processes.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free (ab183216)

This IHC data was generated using the same anti-LRRK2 antibody clone, MJFF3 (c69-6), in a different buffer formulation (cat# **ab133475**).

ab133475, at 1/200 dilution, staining LRRK2 in paraffin embedded Human hippocampus tissue using immunohistochemical analysis.

Heat mediated antigen retrieval was performed with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free (ab183216)

This WB data was generated using the same anti-LRRK2 antibody clone, MJFF3 (c69-6), in a different buffer formulation (cat# **ab133475**).

Lane 1: A549 whole cell lysate (20 µg)


Lane 2: MEF whole cell lysate (20 µg)

Lane 3: A549 (KO) whole cell lysate (20 µg)

Lane 4: MEF (KO) whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - **ab133475** observed at 286 kDa. Red - loading control, **ab18058**, observed at 130 kDa.

ab133475 was shown to specifically react with Wild type A549 and MEF cells whilst no band was observed when knockout samples were used. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and knockout samples were subjected to SDS-PAGE. Ab133475 and **ab18058** (Mouse anti Vinculin loading control) were incubated overnight at 4°C at 1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-LRRK2 antibody [MJFF3 (c69-6)]
- BSA and Azide free (ab183216)

Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free
(ab183216) + NIH/3T3 (mouse embryo) whole cell lysate at 10 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#))

Predicted band size: 286 kDa

Exposure time: 10 seconds

Blocking buffer and concentration: 5% NFDM/TBST

Diluting buffer and concentration: 5% NFDM/TBST

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-LRRK2 antibody [MJFF3 (c69-6)] - BSA and Azide free (ab183216)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors