

Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free ab185129

KO 評価済 リコンビナント RabMAb

画像数 11

製品の概要

製品名	Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR2994(2)] to Lysozyme - BSA and Azide free
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB, ICC/IF 適用なし: IP
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: THP-1, HepG2, RAW 264.7 and HL-60 whole cell lysate; Human spleen tissue lysate; Natural human Lysozyme protein. IHC-P: Human tonsil, spleen, lung, kidney, brain, breast and heart tissues; Mouse spleen and small intestine tissues. ICC/IF: THP-1 cells.
特記事項	<p>ab185129 is the carrier-free version of ab108508.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.20 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリクローナル	モノクローナル
クローン名	EPR2994(2)
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab185129の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. Heat up to 98 degrees C, below boiling, and then let cool for 10-20 min.
WB		Use at an assay dependent concentration. Predicted molecular weight: 17 kDa.
ICC/IF		Use at an assay dependent concentration.

追加情報 Is unsuitable for IP.

ターゲット情報

機能	Lysozymes have primarily a bacteriolytic function; those in tissues and body fluids are associated with the monocyte-macrophage system and enhance the activity of immunogens.
関連疾患	Amyloidosis 8
配列類似性	Belongs to the glycosyl hydrolase 22 family.
細胞内局在	Secreted.



All lanes : Anti-Lysozyme antibody [EPR2994(2)] ([ab108508](#)) at 1/10000 dilution

Lane 1 : Wild-type THP-1 (Human monocytic leukemia cell line) whole cell lysate

Lane 2 : LYZ knockout THP-1 (Human monocytic leukemia cell line) whole cell lysate

Lane 3 : Hep G2 (Human liver hepatocellular carcinoma cell line) whole cell lysate

Lane 4 : HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 17 kDa

Observed band size: 16 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab108508](#)).

Lanes 1 - 4: Merged signal (red and green). Green - [ab108508](#) observed at 16 kDa. Red - loading control [ab8245](#) (Mouse anti-GAPDH antibody [6C5]) observed at 37 kDa.


[ab108508](#) was shown to react with Lysozyme in wild-type THP1 cells in Western blot with loss of signal observed in LYZ knockout sample. Wild-type THP1 and LYZ knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween[®]) before incubation with [ab108508](#) and [ab8245](#) (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 10000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Immunocytochemistry/Immunofluorescence analysis of THP-1 (Human monocytic leukemia) cells labeling lysozyme with purified **ab108508** at 1/250. Cells were fixed with 100% methanol. **ab150077**, Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. Cells were counter-stained with **ab195889** Anti-Alpha Tubulin antibody [DM1A] (1/200, 2.5 g/mL) - Microtubule Marker (Alexa Fluor[®]594) at 1/200. DAPI (blue) was used as a nuclear counterstain. Secondary Only Control: PBS was used instead of the primary antibody as the negative control.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue sections labeling lysozyme with purified **ab108508** at a dilution of 1/1500 (0.6 µg/ml). **ab97051** Goat Anti-Rabbit IgG H&L (HRP) at 1/500 was used as the secondary antibody. Sections were counterstained with hematoxylin. Antigen retrieval was heat mediated using EDTA Buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control and is shown in the inset.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).



Immunohistochemical analysis of paraffin-embedded human lung tissue sections labeling lysozyme with purified **ab108508** at a dilution of 1/1500 (0.6 µg/ml). **ab97051** Goat Anti-Rabbit IgG H&L (HRP) at 1/500 was used as the secondary antibody. Sections were counterstained with Hematoxylin. Antigen retrieval was heat mediated using EDTA Buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control and is shown in the inset.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)




Unpurified **ab108508**, at 1/1000 dilution, staining Lysozyme in Human tonsil by Immunohistochemistry, Paraffin-embedded tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Unpurified **ab108508**, at 1/1000 dilution, staining Lysozyme in Human spleen by Immunohistochemistry, Paraffin-embedded tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Unpurified **ab108508**, at 1/1000 dilution, staining Lysozyme in Human kidney by Immunohistochemistry, Paraffin-embedded tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Unpurified **ab108508** showing negative staining in Normal brain tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Unpurified **ab108508** showing negative staining in Normal breast tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Unpurified **ab108508** showing negative staining in Normal heart tissue.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108508**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Lysozyme antibody [EPR2994(2)] - BSA and Azide free (ab185129)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors