

# Anti-Macrophage Inflammatory Protein 1 gamma antibody ab9913

## 2 References

### 製品の概要

製品名	Anti-Macrophage Inflammatory Protein 1 gamma antibody
製品の詳細	Rabbit polyclonal to Macrophage Inflammatory Protein 1 gamma
由来種	Rabbit
アプリケーション	<b>適用あり:</b> IHC-P, WB, ELISA
種交差性	<b>交差種:</b> Mouse
免疫原	Recombinant full length protein corresponding to Mouse Macrophage Inflammatory Protein 1 gamma. Highly pure (>98%) recombinant mMIP-1g (Mouse macrophage inflammatory protein-1 gamma) Database link: <b><a href="#">P51670</a></b>
ポジティブ・コントロール	Recombinant mouse Macrophage Inflammatory Protein 1 gamma ( <b><a href="#">ab9914</a></b> ) can be used as a positive control in WB.
特記事項	Although some customers have been successful in IHC we no longer batch test in this application.  The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.  If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

### 製品の特性

製品の状態	Lyophilized:Reconstitute with 200µl of sterile water.
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	No preservative, sterile filtered
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	unknown
軽鎖の種類	unknown

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab9913の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. PubMed: 20799334
WB		Use at an assay dependent concentration. To detect mMIP-1g by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mMIP-1g is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
ELISA		Use at an assay dependent concentration. To detect mMIP-1g by direct ELISA (using 100µl/well antibody solution) a concentration of at least 0.5µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant mMIP-1g.

## ターゲット情報

機能	Monokine with inflammatory, pyrogenic and chemokinetic properties. Circulates at high concentrations in the blood of healthy animals. Binding to a high-affinity receptor activates calcium release in neutrophils. It also inhibits colony formation of bone marrow myeloid immature progenitors.
組織特異性	Expressed mainly in the liver, lung, and the thymus, although some expression has been detected in a wide variety of tissues except brain.
配列類似性	Belongs to the intercrine beta (chemokine CC) family.
細胞内局在	Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors