


Anti-MAP2 antibody [HM-2] ab11267

★★★★★ **8 Abreviews** **158 References** 画像数 3

製品の概要

製品名	Anti-MAP2 antibody [HM-2]
製品の詳細	Mouse monoclonal [HM-2] to MAP2
由来種	Mouse
アプリケーション	適用あり: ICC, WB
種交差性	交差種: Rat 交差が予測される動物種: Mouse, Chicken, Cow, Human, Quail 
免疫原	Full length protein corresponding to Rat MAP2. Rat brain microtubule associated proteins (MAPs).
ポジティブ・コントロール	WB: Rat brain tissue lysate. ICC: B35 cells.
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 05/06/2019. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>Storage in frost-free freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: 0.0268% PBS

精製度	Tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	HM-2
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab11267の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC	★★★★★ (2)	Use a concentration of 2 µg/ml. Fix cells in 4% paraformaldehyde/PBS for 45 min; then permeabilise cells with 0.2% Triton X-100 in PBS for 5 min (see Farah et al) OR fix in 4% paraformaldehyde (containing 0.2% picric acid in 0.1 M phosphate buffer, pH 6.9) for 15 min at room temperature (see O' Hare et al) .
WB	★★★★★ (2)	Use a concentration of 1 - 2 µg/ml. Predicted molecular weight: 200 kDa.

ターゲット情報

機能	The exact function of MAP2 is unknown but MAPs may stabilize the microtubules against depolymerization. They also seem to have a stiffening effect on microtubules.
配列類似性	Contains 3 Tau/MAP repeats.
翻訳後修飾	Phosphorylated at serine residues in K-X-G-S motifs by MAP/microtubule affinity-regulating kinase (MARK1 or MARK2), causing detachment from microtubules, and their disassembly (By similarity). Isoform 2 is probably phosphorylated by PKA at Ser-323, Ser-354 and Ser-386 and by FYN at Tyr-67.
細胞内局在	Cytoplasm, cytoskeleton.

画像



Immunocytochemistry - Anti-MAP2 antibody [HM-2]
(ab11267)

Bailey, J.A. et al PLoS One. 2011;6(7):e21954. doi:
10.1371/journal.pone.0021954. Epub 2011 Jul 22
Reproduced under the Creative Commons license
<https://creativecommons.org/publicdomain/zero/1.0/>


Rivastigmine preserves neuronal morphology and alters sAPP secretion

At day 12 or day 20, cells were fixed for immunocytochemistry analysis and probed with anti-MAP2 (green) and anti-GFAP (red) antibodies. Neuronal MAP2 immunoreactivity declines to almost undetectable levels between day 12 and day 20 in untreated cells, whereas neuronal morphology is preserved in rivastigmine treated cultures. Glial GFAP was observed to increase in both treated and untreated cells between day 12 and day 20. These results confirm the degeneration of neurons by day 20.

MAP2 is detected using ab11267 in 4% paraformaldehyde-fixed, 0.5% Triton X-100 permeabilized embryonic rat cerebrocortical cells.

(From Figure 2A of Bailey et al)

This image was generated using the ascites version of the product.




Western blot - Anti-MAP2 antibody [HM-2]
(ab11267)

Anti-MAP2 antibody [HM-2] (ab11267) at 1 µg/ml + Rat brain lysate

Predicted band size: 200 kDa

This image was generated using the ascites version of the product.



Immunocytochemistry - Anti-MAP2 antibody [HM-2]
(ab11267)

Immunocytochemical analysis of B35 (rat neuroblastoma cell line) cells labelling MAP2 with ab11267 at a concentration of 2 µg/mL. The secondary was developed using Goat anti-mouse IgG. Cells were counterstained with DAPI to stain nuclei.

This image was generated using the ascites version of the product.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors