




Anti-MDM2 (phospho S166) antibody ab131355

5 References **画像数 3**

製品の概要

製品名	Anti-MDM2 (phospho S166) antibody
製品の詳細	Rabbit polyclonal to MDM2 (phospho S166)
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF
種交差性	交差種: Human 交差が予測される動物種: Mouse 
免疫原	Synthetic peptide corresponding to Human MDM2 conjugated to keyhole limpet haemocyanin. Synthetic phosphopeptide conjugated to KLH, surrounding phosphorylation site of Serine 166 (A-I-S(p)-E-T)) derived from Human MDM2 (NP_002383.2). Database link: Q00987-1 <div>  Run BLAST with  Run BLAST with </div>
ポジティブ・コントロール	293 cell extract treated with Hydroxyurea, HeLa cells, Human breast carcinoma tissue
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.88% Sodium chloride PBS is without Mg ²⁺ and Ca ²⁺
精製度	Immunogen affinity purified

特記事項 (精製)	ab131355 is purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab131355の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Detects a band of approximately 90 kDa (predicted molecular weight: 55 kDa).
IHC-P		1/50 - 1/100.
ICC/IF		1/100 - 1/200.

ターゲット情報


機能	E3 ubiquitin-protein ligase that mediates ubiquitination of p53/TP53, leading to its degradation by the proteasome. Inhibits p53/TP53- and p73/TP73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Also acts as an ubiquitin ligase E3 toward itself and ARRB1. Permits the nuclear export of p53/TP53. Promotes proteasome-dependent ubiquitin-independent degradation of retinoblastoma RB1 protein. Inhibits DAXX-mediated apoptosis by inducing its ubiquitination and degradation. Component of the TRIM28/KAP1-MDM2-p53/TP53 complex involved in stabilizing p53/TP53. Also component of the TRIM28/KAP1-ERBB4-MDM2 complex which links growth factor and DNA damage response pathways.
組織特異性	Ubiquitous. Isoform Mdm2-A, isoform Mdm2-B, isoform Mdm2-C, isoform Mdm2-D, isoform Mdm2-E, isoform Mdm2-F and isoform Mdm2-G are observed in a range of cancers but absent in normal tissues.
関連疾患	Note=Seems to be amplified in certain tumors (including soft tissue sarcomas, osteosarcomas and gliomas). A higher frequency of splice variants lacking p53 binding domain sequences was found in late-stage and high-grade ovarian and bladder carcinomas. Four of the splice variants show loss of p53 binding.
配列類似性	Belongs to the MDM2/MDM4 family. Contains 1 RanBP2-type zinc finger. Contains 1 RING-type zinc finger. Contains 1 SWIB domain.
ドメイン	Region I is sufficient for binding p53 and inhibiting its G1 arrest and apoptosis functions. It also binds p73 and E2F1. Region II contains most of a central acidic region required for interaction with ribosomal protein L5 and a putative C4-type zinc finger. The RING finger domain which coordinates two molecules of zinc interacts specifically with RNA whether or not zinc is present and mediates the heterooligomerization with MDM4. It is also essential for its ubiquitin ligase E3 activity toward p53 and itself.
翻訳後修飾	Phosphorylated in response to ionizing radiation in an ATM-dependent manner.

Auto-ubiquitinated; which leads to proteasomal degradation. Deubiquitinated by USP2 leads to its accumulation and increases deubiquitination and degradation of p53/TP53. Deubiquitinated by USP7; leading to stabilize it.

細胞内局在

Nucleus > nucleoplasm. Cytoplasm. Nucleus > nucleolus. Expressed predominantly in the nucleoplasm. Interaction with ARF(P14) results in the localization of both proteins to the nucleolus. The nucleolar localization signals in both ARF(P14) and MDM2 may be necessary to allow efficient nucleolar localization of both proteins. Colocalizes with RASSF1 isoform A in the nucleus.

画像



Western blot - Anti-MDM2 (phospho S166) antibody (ab131355)


All lanes : Anti-MDM2 (phospho S166) antibody (ab131355) at 1/500 dilution

Lane 1 : 293 cell extract untreated

Lane 2 : 293 cell extract treated with hydroxyurea


Predicted band size: 55 kDa

Observed band size: 90 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MDM2 (phospho S166) antibody (ab131355)

Immunohistochemical analysis of paraffin embedded Human breast carcinoma tissue labelling MDM2 (phospho S166) with ab131355 at a dilution of 1/50. Picture on the right was preincubated with blocking peptide.



Immunocytochemistry/ Immunofluorescence - Anti-MDM2 (phospho S166) antibody (ab131355)

Immunofluorescence of methanol-fixed HeLa cells staining MDM2 (phospho S166) with ab131355 at a dilution of 1/100.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors