

# Anti-MHC Class II antibody [MRC OX-6] - BSA and Azide free ab237959

リコンビナント

画像数 4

### 製品の概要

製品名	Anti-MHC Class II antibody [MRC OX-6] - BSA and Azide free
製品の詳細	Mouse monoclonal [MRC OX-6] to MHC Class II - BSA and Azide free
由来種	Mouse
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-P
種交差性	<b>交差種:</b> Mouse, Rat
免疫原	Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Rat spleen tissue; IHC-fr: Rat spleen tissue. ICC/IF: Mouse and rat splenocytes
特記事項	<p>ab237959 is the carrier-free version of <b>ab23990</b>.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

## 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at +4°C. Do Not Freeze.
バッファー	Constituent: PBS
キャリア・フリー	はい
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	MRC OX-6
ミエローマ	NS1
アイソタイプ	IgG1

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、**次のテスト済みアプリケーションにおけるab237959の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use at an assay dependent concentration.
IHC-P		1/100.

## ターゲット情報

機能	<p>Binds peptides derived from antigens that access the endocytic route of antigen presenting cells (APC) and presents them on the cell surface for recognition by the CD4 T-cells. The peptide binding cleft accomodates peptides of 10-30 residues. The peptides presented by MHC class II molecules are generated mostly by degradation of proteins that access the endocytic route, where they are processed by lysosomal proteases and other hydrolases. Exogenous antigens that have been endocytosed by the APC are thus readily available for presentation via MHC II molecules, and for this reason this antigen presentation pathway is usually referred to as exogenous. As membrane proteins on their way to degradation in lysosomes as part of their normal turn-over are also contained in the endosomal/lysosomal compartments, exogenous antigens must compete with those derived from endogenous components. Autophagy is also a source of endogenous peptides, autophagosomes constitutively fuse with MHC class II loading compartments. In addition to APCs, other cells of the gastrointestinal tract, such as epithelial cells, express MHC class II molecules and CD74 and act as APCs, which is an unusual trait of the GI tract. To produce a MHC class II molecule that presents an antigen, three MHC class II molecules (heterodimers of an alpha and a beta chain) associate with a CD74 trimer in the ER to form an heterononamer. Soon after the entry of this complex into the endosomal/lysosomal system where antigen processing occurs, CD74 undergoes a sequential degradation by various proteases, including CTSS and CTSL, leaving a small fragment termed CLIP (class-II-associated invariant chain peptide). The removal of CLIP is facilitated by HLA-DM via direct binding to the alpha-beta-</p>
----	---

CLIP complex so that CLIP is released. HLA-DM stabilizes MHC class II molecules until primary high affinity antigenic peptides are bound. The MHC II molecule bound to a peptide is then transported to the cell membrane surface. In B cells, the interaction between HLA-DM and MHC class II molecules is regulated by HLA-DO. Primary dendritic cells (DCs) also to express HLA-DO. Lysosomal microenvironment has been implicated in the regulation of antigen loading into MHC II molecules, increased acidification produces increased proteolysis and efficient peptide loading.

#### 配列類似性


Belongs to the MHC class II family.

Contains 1 Ig-like C1-type (immunoglobulin-like) domain.

#### 細胞内局在

Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus > trans-Golgi network membrane. Endosome membrane. Lysosome membrane. The MHC class II complex transits through a number of intracellular compartments in the endocytic pathway until it reaches the cell membrane for antigen presentation.

#### 画像



This data was developed using the same antibody clone in a different buffer formulation containing PBS, L-arginine, and sodium azide ([ab23990](#)).


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Rat splenocyte cells labelling MHC Class II with [ab23990](#) at 1/100 dilution (10.63 ug/ml), followed by [ab150117](#) Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). [ab206369](#) Anti-beta Tubulin rabbit monoclonal antibody (Alexa Fluor® 594) was used at 1/100 dilution (5µg/mL) as counterstain for tubulin (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is [ab150117](#) Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody.

Confocal image showing membranous and cytoplasmic staining in subsets of rat splenocyte.

**Negative control:** C6.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-MHC Class II antibody [MRC OX-6] - BSA and Azide free (ab237959)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, L-arginine, and sodium azide (**ab23990**).

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse splenocyte cells labelling MHC Class II with **ab23990** at 1/100 dilution (10.63 ug/ml), followed by **ab150117** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). **ab206369** Anti-beta Tubulin rabbit monoclonal antibody (Alexa Fluor® 594) was used at 1/100 dilution (5µg/mL) as counterstain for tubulin (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is **ab150117** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) preadsorbed antibody.

Confocal image showing membranous and cytoplasmic staining in subsets of mouse splenocyte . Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MHC Class II antibody [MRC OX-6] - BSA and Azide free (ab237959)

IHC image of MHC Class II staining in Rat normal spleen formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with **ab23990**, 10µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, L-arginine, and sodium azide (**ab23990**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MHC Class II antibody [MRC OX-6] - BSA and Azide free (ab237959)

This image is courtesy of an abreview submitted by Carl Hobbs, King's College London, United Kingdom

Immunohistochemical detection of MHC Class II using antibody **ab23990** on PFA-fixed rat spleen tissue sections. Antibody diluted at 1/100 and incubated for 2 hours in TBS/BSA/Tween/azide. Secondary antibody: anti mouse IgG conjugated to biotin (1/100). After dissection of spleen from PFA-perfused specimen it was sampled and further immersion-fixed for two Hrs. After subsequent immersion in 30% Sucrose, specimens were snap-frozen. Before immunostaining, the 8 micron sections were placed in a 60 degree C oven for 60 mins to enhance adhesion. The submitted image shows white pulp (PALS) and a small area of red pulp (RP-upper left). The Periarterial Sheath (PALS) with its Central Arteriole/artery (CA) shows many positive cells (B-lymphocytes and Macrophages) and negative lymphocytes (T-cells?). The Marginal Sinus (MS) is clearly seen between the PALS and the Marginal Zone (MZ). There is a clear Germinal Centre (GS) in this image.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, L-arginine, and sodium azide (**ab23990**).

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors