# abcam

# Product datasheet

# Mouse IgG2b, kappa monoclonal [MG2b-57] - Isotype control ab18421

# 4 References

#### 製品の概要

製品名 Mouse IgG2b, kappa monoclonal [MG2b-57] - Isotype control

特異性 The MG2b-57 immunoglobulin (ab18421) reacts with KLH. The MG2b-57 immunoglobulin was

chosen as an isotype control after screening on a variety of resting, activated, live, and fixed

mouse and human tissues.

アプリケーション 適用あり: IP, WB, IHC-Fr, IHC-P, Flow Cyt

免疫原 Trinitrophenol + KLH

特記事項 ab18421 MG2b-57 immunoglobulin is useful as an isotype-matched control in Western blotting,

immunoprecipitation, immunohistochemical staining, bioassay and immunofluorescent staining

(surface or intracellular) for flow cytometric analysis.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

**バッファー** pH: 7.20

Preservative: 0.09% Sodium azide

Constituent: PBS

精製度 Immunogen affinity purified

特記事項(精製) The immunoglobulin was purified by affinity chromatography.

アイソタイプ・コントロール 備考 ab18421 MG2b-57 immunoglobulin is useful as an isotype-matched control in Western blotting,

immunoprecipitation, immunohistochemical staining, bioassay and immunofluorescent staining

(surface or intracellular) for flow cytometric analysis.

1

ポリ/モノ モノクローナル クローン名 MG2b-57 アイソタイプ lgG2b 軽鎖の種類 kappa

## アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab18421の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. Surface or intracellular staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors