



Anti-NFAT1 antibody [25A10.D6.D2] ab2722

★★★★★ [7 Abreviews](#) [53 References](#) [画像数 10](#)

製品の概要

製品名	Anti-NFAT1 antibody [25A10.D6.D2]
製品の詳細	Mouse monoclonal [25A10.D6.D2] to NFAT1
由来種	Mouse
特異性	Ab2722 detects nuclear factor of activated T-cells (NFAT) from mouse, rat and human tissues (endogenously expressed). This antibody does not cross react with NFAT2 (NFATc, NFATc1). This antibody detects both forms NFAT1 - a ~140 kDa protein representing phosphorylated NFAT1 in resting immune cells, and a ~120 kDa protein in stimulated cells that represents fully-dephosphorylated NFAT1.
アプリケーション	適用あり: Flow Cyt, WB, IHC-P, ICC/IF
種交差性	交差種: Human
免疫原	Synthetic peptide corresponding to Mouse NFAT1 aa 50-150. Database link: Q60591
	 Run BLAST with  Run BLAST with
ポジティブ・コントロール	WB: resting immune cells and ionomycin stimulated immune cells (see Shaw et al reference: "1uM for T cells and B cells, 10uM for macrophages, and 0.3uM for mast cells. Treatment was for 20 min.")
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.05% Sodium azide Constituent: PBS

精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	25A10.D6.D2
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab2722の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (4)	Use a concentration of 1 µg/ml. Detects a band of approximately 120, 140 kDa (predicted molecular weight: 115 kDa).
IHC-P	★★★★★ (1)	1/50 - 1/200.
ICC/IF		1/100 - 1/1000.

ターゲット情報


機能	Plays a role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2, IL-3, IL-4, TNF-alpha or GM-CSF.
組織特異性	Expressed in thymus, spleen, heart, testis, brain, placenta, muscle and pancreas.
配列類似性	Contains 1 RHD (Rel-like) domain.
ドメイン	Rel Similarity Domain (RSD) allows DNA-binding and cooperative interactions with AP1 factors.
翻訳後修飾	In resting cells, phosphorylated by NFATC-kinase on at least 18 sites in the 99-363 region. Upon cell stimulation, all these sites except Ser-243 are dephosphorylated by calcineurin. Dephosphorylation induces a conformational change that simultaneously exposes an NLS and masks an NES, which results in nuclear localization. Simultaneously, Ser-53 or Ser-56 is phosphorylated; which is required for full transcriptional activity.
細胞内局在	Cytoplasm. Nucleus. Cytoplasmic for the phosphorylated form and nuclear after activation that is controlled by calcineurin-mediated dephosphorylation. Rapid nuclear exit of NFATC is thought to be one mechanism by which cells distinguish between sustained and transient calcium signals. The subcellular localization of NFATC plays a key role in the regulation of gene transcription.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human spleen tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a mouse monoclonal antibody recognizing NFATc2 ab2722 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunocytochemistry/ Immunofluorescence - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunocytochemistry/Immunofluorescence analysis of NFAT1 (green) in HeLa cells. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% BSA for 15 minutes at room temperature. Cells were left untreated (left panel) or treated with 1uM staurosporine (right panel) for 3 hours and incubated with ab2722 (1:100) for at least 1 hour at room temperature, washed with PBS, and incubated with a DyLight 488 conjugated goat anti-mouse IgG secondary antibody (1:400) for 30 minutes at room temperature. F-Actin (red) was stained with DyLight 554 Phalloidin and nuclei (blue) were stained with Hoechst 33342 dye. Images were taken at 20X magnification.



Western blot - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Anti-NFAT1 antibody [25A10.D6.D2] (ab2722) at 1 µg/ml + Human spleen tissue lysate - total protein ([ab29699](#)) at 10 µg

Secondary

Goat Anti-Mouse IgG H&L (HRP) preadsorbed ([ab97040](#)) at 1/5000 dilution ()

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 115 kDa

Observed band size: 150 kDa

Additional bands at: 62 kDa. We are unsure as to the identity of these extra bands.


Exposure time: 4 minutes

NFAT1 contains an extensive number of potential phosphorylation sites (SwissProt) which may explain its migration at a higher molecular weight than predicted.




Flow Cytometry - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Overlay histogram showing Jurkat cells stained with ab2722 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab2722, 2µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

ab2722 (4µg/ml) staining NFAT in human tonsil, using an automated system (DAKO Autostainer Plus). Using this protocol there is strong nuclear and weak cytoplasmic staining. Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer EDTA pH 9.0 in a DAKO PT link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.




Immunocytochemistry/ Immunofluorescence - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunocytochemistry/Immunofluorescence analysis of NFAT1 (green) shows staining in HeLa cells. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2722 (1:20) over night at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.




Immunocytochemistry/ Immunofluorescence - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunocytochemistry/Immunofluorescence analysis of NFAT1 (green) shows staining in MCF-7 cells. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2722 (1:20) overnight at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)


Immunocytochemistry/Immunofluorescence analysis of NFAT1 (green) shows staining in U251 Cells. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2722 (1:20) overnight at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human colon carcinoma tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a mouse monoclonal antibody recognizing NFATc2 ab2722 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase

activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NFAT1 antibody [25A10.D6.D2] (ab2722)

Immunohistochemistry was performed on both normal and cancer biopsies of deparaffinized Human tonsil tissue tissues. To expose target proteins heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature. Tissues were then probed at a dilution of 1:100 with a mouse monoclonal antibody recognizing NFATc2 ab2722 or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors