

Anti-p18 INK4c/CDKN2C antibody [EPR15891] ab192239

KO 評価済 リコンビナント RabMAb

14 References 画像数 8

製品の概要

製品名	Anti-p18 INK4c/CDKN2C antibody [EPR15891]
製品の詳細	Rabbit monoclonal [EPR15891] to p18 INK4c/CDKN2C
由来種	Rabbit
アプリケーション	適用あり: WB, IP, ICC/IF, IHC-P
種交差性	交差種: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HEL-293T, Ramos, HeLa, Jurkat, Rat kidney, Rat spleen and NIH 3T3 lysates. IHC-P: Human brain and glioma tissues; ICC/IF: HeLa cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.05% BSA</p>
精製度	Ion Exchange Chromatography
ポリ/モノ	モノクローナル
クローン名	EPR15891
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab192239の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000. Detects a band of approximately 18 kDa (predicted molecular weight: 18 kDa).
IP		1/30.
ICC/IF		1/100.
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

ターゲット情報

機能	Interacts strongly with CDK6, weakly with CDK4. Inhibits cell growth and proliferation with a correlated dependence on endogenous retinoblastoma protein RB.
組織特異性	Highest levels found in skeletal muscle. Also found in pancreas and heart.
配列類似性	Belongs to the CDKN2 cyclin-dependent kinase inhibitor family. Contains 4 ANK repeats.

画像



All lanes : Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CDKN2C knockout HeLa cell lysate

Lane 3 : Rat Kidney cell lysate

Lane 4 : Jurkat cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 18 kDa

Lanes 1-4: Merged signal (red and green). Green - ab192239


observed at 18 kDa. Red - loading control **ab8245** observed at 37 kDa.

ab192239 Anti-p18 INK4c/CDKN2C antibody [EPR15891] was shown to specifically react with Cyclin Dependent Kinase Inhibitor 2C in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265031** (knockout cell lysate **ab257887**) was used. Wild-type and Cyclin Dependent Kinase Inhibitor 2C knockout samples were subjected to SDS-PAGE. ab192239 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunofluorescent analysis of HeLa cells (4% Paraformaldehyde-fixed, 0.1% tritonX-100 permeabilized) labeling p18 INK4C/CDKN2C with ab192239 at 1/100 dilution (5µg/mL) followed by Goat anti rabbit IgG (AlexaFluor® 488) (**ab150077**) secondary at 1/200 dilution and counter-stained with DAPI (blue). Negative controls: anti-p18 INK4C/CDKN2C at 1/100 dilution, Secondary ab (Goat anti mouse IgG (Alexa Fluor®594)) at 1/400 dilution.

Immunocytochemistry/ Immunofluorescence - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)



Western blot - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)

All lanes : Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239) at 1/1000 dilution

Lane 1 : Rat kidney lysate

Lane 2 : Rat spleen lysate


Lane 3 : NIH 3T3 lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution


Predicted band size: 18 kDa



Immunoprecipitation - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)

Western blot analysis of immunoprecipitation pellet from HeLa lysate immunoprecipitated using ab192239 at 1/30 dilution.


Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)

Immunohistochemical analysis of paraffin-embedded Human glioma tissue labeling p18 INK4C/CDKN2C with ab192239 at 1/50 dilution followed by pre-diluted HRP Polymer for Rabbit/Mouse IgG secondary antibody and counter-stained with Hematoxylin. (inset: negative control).


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemical analysis of paraffin-embedded Human brain tissue labeling p18 INK4C/CDKN2C with ab192239 at 1/50 dilution followed by pre-diluted HRP Polymer for Rabbit/Mouse IgG secondary antibody and counter-stained with Hematoxylin. (inset: negative control).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)



Western blot - Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239)

All lanes : Anti-p18 INK4c/CDKN2C antibody [EPR15891] (ab192239) at 1/10000 dilution

Lane 1 : 293 cell lysate

Lane 2 : Ramos cell lysate

Lane 3 : HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

Predicted band size: 18 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-p18 INK4c/CDKN2C antibody [EPR15891]
(ab192239)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors