

Anti-p21 antibody [EPR3993] ab109199

KO 評価済 リコンビナント RabMAb

★★★★☆ 12 Abreviews 276 References 画像数 11

製品の概要

製品名	Anti-p21 antibody [EPR3993]
製品の詳細	Rabbit monoclonal [EPR3993] to p21
由来種	Rabbit
特異性	<p>Expression levels of the target protein vary between different tissue/cell lines and in some cases, induction may be required before a signal is observed.</p> <p>This antibody is not recommended for use in WB with tissue and primary cell samples.</p> <p>We recommended ab109520 and ab188224 for use in IHC.</p>
アプリケーション	適用あり: WB
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Raw 264.7, HCT116, MCF-7, PC-12 treated with 50ng/ml NFG for 48 hours whole cell lysate, wild-type HeLa Treated Fluvastatin (50 uM, 24 h) cell lysate, Wild-type DLD-1 20 μM 2,3-DCPE for 16hrs treated cell lysate
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol, 59% PBS, 0.05% BSA</p>

精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR3993
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab109199の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (6)	1/1000. Detects a band of approximately 21 kDa (predicted molecular weight: 18 kDa). For unpurified use at 1/1000 - 1/10000. Not recommended for use with tissue samples.

ターゲット情報

機能	May be the important intermediate by which p53/TP53 mediates its role as an inhibitor of cellular proliferation in response to DNA damage. Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex.
組織特異性	Expressed in all adult human tissues, with 5-fold lower levels observed in the brain.
配列類似性	Belongs to the CDI family.
ドメイン	The PIP-box K+4 motif mediates both the interaction with PCNA and the recruitment of the DCX(DTL) complex: while the PIP-box interacts with PCNA, the presence of the K+4 submotif, recruits the DCX(DTL) complex, leading to its ubiquitination. The C-terminal is required for nuclear localization of the cyclin D-CDK4 complex.
翻訳後修飾	Phosphorylation of Thr-145 by Akt or of Ser-146 by PKC impairs binding to PCNA. Phosphorylation at Ser-114 by GSK3-beta enhances ubiquitination by the DCX(DTL) complex. Ubiquitinated by MKRN1; leading to polyubiquitination and 26S proteasome-dependent degradation. Ubiquitinated by the DCX(DTL) complex, also named CRL4(CDT2) complex, leading to its degradation during S phase or following UV irradiation. Ubiquitination by the DCX(DTL) complex is essential to control replication licensing and is PCNA-dependent: interacts with PCNA via its PIP-box, while the presence of the containing the 'K+4' motif in the PIP box, recruit the DCX(DTL) complex, leading to its degradation.
細胞内局在	Cytoplasm. Nucleus.

画像



All lanes : Anti-p21 antibody [EPR3993] (ab109199) at 1/1000 dilution

Lane 1 : Raw 264.7(Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lane 3 : C6 (Rat glial tumor glial cell) whole cell lysate

Lane 4 : PC-12(Rat adrenal gland pheochromocytoma) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 18 kDa

Observed band size: 18 kDa

Exposure time: 180 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

[ab181602](#) was used as a GAPDH loading control.



All lanes : Anti-p21 antibody [EPR3993] (ab109199) at 1/1000 dilution

Lane 1 : PC-12(Rat adrenal gland pheochromocytoma) whole cell lysate

Lane 2 : PC-12(Rat adrenal gland pheochromocytoma) treated with 50ng/ml NFG for 48 hours whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Developed using the ECL technique.

Predicted band size: 18 kDa

Observed band size: 18 kDa

Exposure time: 180 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

ab181602 was used as a GAPDH loading control.

We recommend using higher or super higher sensitivity ECL substrate for detecting.

Increase lysate amount can also help to get stronger signal.

All lanes : Anti-p21 antibody [EPR3993] (ab109199) at 1/1000 dilution

Lane 1 : wild-type HeLa Vehicle Control Fluvastatin (0 uM, 24 h) cell lysate

Lane 2 : wild-type HeLa Treated Fluvastatin (50 uM, 24 h) cell lysate

Lane 3 : CDKN1A knockout HeLa Vehicle Control Fluvastatin (0 uM, 24 h) cell lysate


Lane 4 : CDKN1A knockout HeLa Treated Fluvastatin (50 uM, 24 h) cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 18 kDa

Observed band size: 21 kDa



False colour image of Western blot: Anti-p21 antibody [EPR3993] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (**ab7291**) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab109199 was shown to bind specifically to p21. A band was observed at 21 kDa in wild-type HeLa cell lysates with no signal observed at this size in CDKN2A knockout cell line **ab255349** (knockout cell lysate **ab263812**). To generate this image, wild-type and CDKN2A knockout cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were

blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Lane 1: Wild-type DLD-1 cell lysate (20 µg)

Lane 2: Wild-type DLD-1 20 µM 2,3-DCPE for 16hrs treated cell lysate (20 µg)


Lane 3: p21 knockout DLD-1 cell lysate (20 µg)

Lane 4: p21 knockout 20 µM 2,3-DCPE for 16hrs DLD-1 cell lysate (20 µg)

Lane 5: HT1080 cell lysate (20 µg)

Lanes 1 - 5: Merged signal (red and green). Green - ab109199 observed at 20 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab109199 was shown to recognize p21 in WT DLD-1 cells with 2,3-DCPE treatment along with additional cross-reactive bands. When p21 knockout DLD-1 cells +/- 2,3-DCPE treatment were used, no band was observed. Wild-type and p21 knockout samples were subjected to SDS-PAGE. ab109199 and [ab8245](#) (loading control to GAPDH) were diluted 1/1000 and 1/10 000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-p21 antibody [EPR3993]
(ab109199)

All lanes : Anti-p21 antibody [EPR3993] (ab109199) at 1/1000 dilution

Lane 1 : Wild-type HCT116 cell lysate

Lane 2 : CDKN1A knockout HCT116 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 18 kDa

Observed band size: 20 kDa


Lanes 1- 2: Merged signal (red and green). Green - ab109199 observed at 20 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

ab109199 was shown to react with p21 in wild-type HCT116 cells in western blot. Loss of signal was observed when knockout cell line [ab266860](#) (knockout cell lysate [ab256870](#)) was used. Wild-Type HCT116 and CDKN1A knockout HCT116 cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab109199 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-p21 antibody [EPR3993]
(ab109199)

Different batches of ab109199 were tested on MCF7 (Human breast adenocarcinoma epithelial cell) lysate at 0.2 µg/ml. 15 µg of lysate was loaded in each lane. Bands observed at 21 kDa.



Western blot - Anti-p21 antibody [EPR3993]
(ab109199)

Lane 1 : Anti-p21 antibody [EPR3993] (ab109199) (0.7ug/ul)

Lane 2 : Anti-p21 antibody [EPR362] (**ab109520**) (0.8ug/ul)

All lanes : MCF-7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

Lysates/proteins at 15 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 18 kDa

Exposure time: 3 minutes

Blocking and diluting buffer: 5% NDFM/TBST.



Lane 1 : Anti-p21 antibody [EPR3993] (ab109199) (1.4ug/ul)

Lane 2 : Anti-p21 antibody [EPR18021] (**ab188224**) (1.0ug/ul)

Lane 1 : RAW264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates


Lane 2 : Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysates

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 18 kDa



Lane 1: Wild-type DLD-1 cell lysate (20 µg)

Lane 2: Wild-type DLD-1 20 µM 2,3-DCPE for 16hrs treated cell lysate (20 µg)


Lane 3: p21 knockout DLD-1 cell lysate (20 µg)

Lane 4: p21 knockout 20 µM 2,3-DCPE for 16hrs DLD-1 cell lysate (20 µg)

Lane 5: HT1080 cell lysate (20 µg)

Lanes 1 - 5: Merged signal (red and green). Green - ab109119 observed at 20 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

This western blot image is a comparison between **ab109119** and a competitor's top cited rabbit polyclonal antibody.



Western blot - Anti-p21 antibody [EPR3993]
(ab109199)

Anti-p21 antibody [EPR3993] (ab109199) at 1/1000 dilution
(purified) + PC-12 cell lysate at 10 µg

Secondary

Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 18 kDa

Observed band size: 21 kDa

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-p21 antibody [EPR3993] (ab109199)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors