

### Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] ab40795

リコンビナント **RabMAb**

★★★★★ **4 Abreviews** **35 References** 画像数 **13**

#### 製品の概要

製品名	Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y]
製品の詳細	Rabbit monoclonal [EP656Y] to PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154)
由来種	Rabbit
アプリケーション	<b>適用あり:</b> Flow Cyt (Intra), WB, IHC-P, IP, ICC/IF
種交差性	<b>交差種:</b> Mouse, Rat, Human
免疫原	Synthetic peptide within Human PAK1 (phospho S144). The exact sequence is proprietary. Database link: <a href="#">Q13153</a>
ポジティブ・コントロール	WB: MCF7, HeLa, RAW 264.7 and C6 cell lysates. IHC: Human liver carcinoma, mouse cerebral cortex, rat cerebral cortex. ICC/IF: HeLa cells. IP: HeLa cell lysate. Flow Cyt (intra): NIH/3T3 cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS</p>

精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP656Y
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee      Abpromise保証は、次のテスト済みアプリケーションにおけるab40795の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/120. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (2)	1/10000 - 1/50000. Detects a band of approximately 66 kDa (predicted molecular weight: 65 kDa).
IHC-P	★★★★☆ (1)	1/100 - 1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/40.
ICC/IF	★★★★★ (1)	1/250 - 1/500.

ターゲット情報

細胞内局在      PAK1: Cytoplasm. Cell junction > focal adhesion. Recruited to focal adhesions upon activation.  
 PAK2: Cytoplasm and Nucleus. Cytoplasm > perinuclear region. Membrane. Interaction with ARHGAP10 probably changes PAK-2p34 location to cytoplasmic perinuclear region.  
 Myristoylation changes PAK-2p34 location to the membrane. PAK3: Cytoplasmic

画像



Western blot - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

**All lanes :** Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795) at 1/1000 dilution

**Lane 1 :** MCF7, grown in serum-free media overnight, whole cell lysate

**Lane 2 :** MCF7, grown in serum-free media overnight, then treated with EGF 1µg/ml for 10min, whole cell lysate

**Lane 3 :** MCF7, grown in serum-free media overnight, then treated with EGF 1µg/ml for 10min, whole cell lysate. The membrane was incubated with phosphatase.

Lysates/proteins at 10 µg per lane.

### Secondary


**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 65 kDa

**Observed band size:** 55 kDa

**Exposure time:** 1 minute


Blocking and dilution buffer: 5% NFDM/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

ab40795 staining PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) in rat cerebral cortex tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) **ab97051** was used as the secondary antibody at a dilution of 1/500.

**Negative control 1:** PBS in place of primary




Immunocytochemistry/ Immunofluorescence - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

ab40795 staining PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) in HeLa (human cervix adenocarcinoma) cells, treated and untreated with Lambda Protein Phosphatase 31I for 5h by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti rabbit IgG (Alexa Fluor® 488) (**ab150077**) was used as the secondary antibody. **ab7291** and **ab150120** were used as counterstains for primary antibody **ab75748** and secondary antibody **ab150077** respectively and DAPI was used as a nuclear counterstain.

**Negative control 1:** Rabbit primary antibody and anti-mouse secondary antibody (**ab150120**)

**Negative control 2:** Mouse primary antibody (**ab7291**) and anti-rabbit secondary antibody (**ab150077**)




Flow Cytometry (Intracellular) - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

ab40795 staining PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) in the human cell line NIH/3T3 (mouse embryo) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/120. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/500 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)




Immunoprecipitation - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

ab40795 immunoprecipitating PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154). 10µg of HeLa (human cervix adenocarcinoma) whole cell lysate was incubated with primary antibody at a dilution of 1/40 and VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) at a dilution of 1/10000.

**Lane 1:** HeLa whole cell lysate (10ug)

**Lane 2:** ab40795 IP in HeLa whole cell lysate

**Lane 3:** Rabbit monoclonal IgG ([ab172730](#)) instead of ab40795 in HeLa (human cervix adenocarcinoma) whole cell lysate



**All lanes :** Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795) at 1/2000 dilution

**Lane 1 :** HeLa cell lysate with None


**Lane 2 :** HeLa cell lysate with PAK2 (pS141)

**Lane 3 :** HeLa cell lysate with PAK2 non-phospho

**Lane 4 :** HeLa cell lysate with PAK3 (pS154)

**Lane 5 :** HeLa cell lysate with PAK3 non-phospho

**Predicted band size:** 65 kDa



**All lanes :** Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795) at 1/50000 dilution

**Lane 1 :** C6 (rat glioma) whole cell lysate - treated with phosphatase


**Lane 2 :** C6 (rat glioma) whole cell lysate - untreated

Lysates/proteins at 10 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


**Predicted band size:** 65 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

ab40795 staining PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) in mouse cerebral cortex tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) [ab97051](#) was used as the secondary antibody at a dilution of 1/500.


**Negative control 1:** PBS in place of primary antibody.



Flow Cytometry (Intracellular) - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

Overlay histogram showing HeLa cells stained with unpurified ab40795 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab40795, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.





**All lanes :** Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795) at 1/10000 dilution

**Lane 1 :** HeLa whole cell lysate - untreated


**Lane 2 :** HeLa whole cell lysate - treated with phosphatase

Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 65 kDa




ab40795 staining PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) in human liver carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with paraformaldehyde and antigen retrieval was by heat mediation in a EDTA buffer. Samples were incubated with primary antibody at a dilution of 1/100. A goat anti-rabbit IgG H&L (HRP) [ab97051](#) was used as the secondary antibody at a dilution of 1/500.

**Negative control 1:** PBS in place of primary antibody.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)





**All lanes :** Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795) at 1/10000 dilution

**Lane 1 :** RAW264.7 (mouse abelson murine leukemia virus-induced tumor) whole cell lysate - treated with phosphatase

**Lane 2 :** RAW264.7 (mouse abelson murine leukemia virus-induced tumor) whole cell lysate - untreated

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

**Predicted band size:** 65 kDa

Why choose a recombinant antibody?

**Research with confidence**  
Consistent and reproducible results

**Long-term and scalable supply**  
Recombinant technology

**Success from the first experiment**  
Confirmed specificity

**Ethical standards compliant**  
Animal-free production

Anti-PAK1 (phospho S144) + PAK2 (phospho S141) + PAK3 (phospho S154) antibody [EP656Y] (ab40795)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors