

Anti-PAK2 antibody [EP796Y] ab76293

KO 評価済 リコンビナント RabMAb

★★★★☆ 2 Abreviews 12 References 画像数 14

製品の概要

製品名	Anti-PAK2 antibody [EP796Y]
製品の詳細	Rabbit monoclonal [EP796Y] to PAK2
由来種	Rabbit
アプリケーション	適用あり: IP, ICC/IF, WB, IHC-P, Flow Cyt (Intra)
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide within Human PAK2 aa 1-100 (N terminal). The exact sequence is proprietary. Database link: Q13177
ポジティブ・コントロール	WB: HeLa, NIH/3T3, RAW 264.7, Wild-type HEK-293T, PAK2 CRISPR-Cas9 edited HEK-293T and C6 cell lysates. IHC-P; Human breast carcinoma tissue IF/ICC: T47D cell line. Flow Cyt (intra): HeLa cells. IP: HeLa and NIH/3T3 cell lysates.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP796Y

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab76293の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IP		1/30.
ICC/IF	★★★★★ (1)	1/100 - 1/250.
WB	★★★★★ (1)	1/5000. Predicted molecular weight: 58 kDa. For unpurified use at 1/1000 - 1/2000.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. This antibody may not be suitable for IHC with mouse or rat samples Use of HRP conjugated or polymerized HRP secondary antibody is recommended. Stronger signals have been found using the polymerized HRP secondary.
Flow Cyt (Intra)		1/20. For unpurified use at 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.


ターゲット情報

機能	The activated kinase acts on a variety of targets. Phosphorylates ribosomal protein S6, histone H4 and myelin basic protein. Full length PAK 2 stimulates cell survival and cell growth. The process is, at least in part, mediated by phosphorylation and inhibition of pro-apoptotic BAD. Caspase-activated PAK-2p34 is involved in cell death response, probably involving the JNK signaling pathway. Cleaved PAK-2p34 seems to have a higher activity than the CDC42-activated form.
組織特異性	Ubiquitously expressed. Higher levels seen in skeletal muscle, ovary, thymus and spleen.
配列類似性	Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily. Contains 1 CRIB domain. Contains 1 protein kinase domain.
翻訳後修飾	Full length PAK 2 is autophosphorylated when activated by CDC42/p21. Following cleavage, both peptides, PAK-2p27 and PAK-2p34, become highly autophosphorylated, with PAK-2p27 being phosphorylated on serine and PAK-2p34 on threonine residues, respectively. Autophosphorylation of PAK-2p27 can occur in the absence of any effectors and is dependent on phosphorylation of Thr-402, because PAK-2p27 is acting as an exogenous substrate. During apoptosis proteolytically cleaved by caspase-3 or caspase-3-like proteases to yield active PAK-2p34. Ubiquitinated, leading to its proteasomal degradation. PAK-2p34 is myristoylated.

細胞内局在

Cytoplasm and Nucleus. Cytoplasm > perinuclear region. Membrane. Interaction with ARHGAP10 probably changes PAK-2p34 location to cytoplasmic perinuclear region. Myristoylation changes PAK-2p34 location to the membrane.

画像



All lanes : Anti-PAK2 antibody [EP796Y] (ab76293) at 1/5000 dilution

Lane 1 : Wild-type HEK-293T cell lysate

Lane 2 : PAK2 CRISPR-Cas9 edited HEK-293T cell lysate

Lane 3 : Wild-type HeLa [ab255552](#) cell lysate

Lane 4 : PAK2 knockout HeLa [ab260287](#) cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution


Performed under reducing conditions.

Predicted band size: 58 kDa

Observed band size: 65 kDa

False colour image of Western blot: Anti-PAK2 antibody [EP796Y] staining at 1/5000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab76293 was shown to bind specifically to PAK2. A band was observed at 65 kDa in wild-type HEK-293T cell lysates with no signal observed at this size in PAK2 CRISPR-Cas9 edited cell line [ab282648](#) (CRISPR-Cas9 edited cell lysate [ab283047](#)). The band observed in the CRISPR-Cas9 edited lysate lane below 65 kDa is likely to represent a truncated form of PAK2. This has not been investigated further and the functional properties of the gene product have not been determined. To generate this image, wild-type and PAK2 CRISPR-Cas9 edited HEK-293T cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary

antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Western blot - Anti-PAK2 antibody [EP796Y] (ab76293)

All lanes : Anti-PAK2 antibody [EP796Y] (ab76293) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : PAK2 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 58 kDa

Observed band size: 60 kDa

Lanes 1- 2: Merged signal (red and green). Green - ab76293 observed at 60 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

ab76293 was shown to react with PAK2 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line [ab264814](#) (knockout cell lysate [ab257573](#)) was used. Wild-type HeLa and PAK2 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab76293 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



PAK2 was immunoprecipitated from 0.35 mg NIH/3T3 (Mouse embryonic fibroblast) cell lysate 10 µg with ab76293 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab76293 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)([ab131366](#)) was used at 1/5000 dilution.


Lane 1: NIH/3T3 (Mouse embryonic fibroblast) cell lysate 10 µg

Lane 2: ab76293 IP in NIH/3T3 cell lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab76293 in HeLa cell lysate


Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 7 seconds




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue sections labeling PAK2 with Purified ab76293 at 1:100 dilution (2.02 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAK2 antibody [EP796Y]
(ab76293)




Immunocytochemistry/ Immunofluorescence - Anti-PAK2 antibody [EP796Y] (ab76293)

Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling PAK2 with purified ab76293 at 1:100 dilution (2.0µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). **ab150077** Goat anti rabbit IgG(Alexa Fluor® 488) was used as the secondary antibody at 1:1000 dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-PAK2 antibody [EP796Y] (ab76293)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling PAK2 with purified ab76293 at 1/20 dilution (10 ug/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilized with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: PAK2 knockout HAP1 cell lysate (20 µg)

Lane 3: HeLa cell lysate (20 µg)

Lane 4: Human lymph node tissue lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab76293 observed at 60 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

Unpurified ab76293 was shown to specifically react with PAK2 when PAK2 knockout samples were used. Wild-type and PAK2 knockout samples were subjected to SDS-PAGE. ab76293 and **ab8245** (loading control to GAPDH) were diluted 1/1000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



All lanes : Anti-PAK2 antibody [EP796Y] (ab76293) at 1/5000 dilution (purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates

Lane 3 : RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates

Lane 4 : C6 (Rat glial tumor glial cell) whole cell lysates


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/2000 dilution

Predicted band size: 58 kDa

Blocking and diluting buffer: 5% NFDM/TBST




Anti-PAK2 antibody [EP796Y] (ab76293) at 1/2000 dilution
(unpurified) + HeLa cell lysate at 10 µg

Secondary

HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 58 kDa

Observed band size: 61 kDa



PAK2 was immunoprecipitated from 0.35 mg HeLa (Human cervix adenocarcinoma epithelial cell) cell lysate 10 µg with ab76293 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab76293 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)([ab131366](#)) was used at 1/5000 dilution.


Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) cell lysate 10 µg

Lane 2: ab76293 IP in HeLa cell lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab76293 in HeLa cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 7 seconds



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PAK2 antibody [EP796Y] (ab76293)


Unpurified ab76293, at a 1/100 dilution, staining PAK2 in paraffin embedded human breast carcinoma tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-PAK2 antibody [EP796Y] (ab76293)

ICC/IF image of unpurified ab76293 stained T47D cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab76293, 1µg/ml) overnight at +4°C. The secondary antibody (green) was **ab96899**, DyLight® 488 goat anti-rabbit IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Flow Cytometry (Intracellular) - Anti-PAK2 antibody [EP796Y] (ab76293)

Overlay histogram showing HeLa cells stained with unpurified ab76293 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab76293, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PAK2 antibody [EP796Y] (ab76293)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors