

PE Anti-beta 2 Microglobulin antibody [B2M-01] ab49424

2 References

製品の概要

製品名	PE Anti-beta 2 Microglobulin antibody [B2M-01]
製品の詳細	PE Mouse monoclonal [B2M-01] to beta 2 Microglobulin
由来種	Mouse
標識	PE. Ex: 488nm, Em: 575nm
特異性	This antibody reacts with beta 2 microglobulin (b2M) associated with cell-surface MHC Class I molecules and other membrane antigens as well as with soluble beta 2 microglobulin. Beta 2 microglobulin is a 12 kDa Ig like glycoprotein expressed on lymphocytes, thymocytes, monocytes, granulocytes, platelets, endothelial cells and epithelial cells. It is absent on erythrocytes.
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Human 非交差種: Mouse, Rabbit, Chicken, Cow, Dog
免疫原	Full length native protein (purified) (Human)
ポジティブ・コントロール	Peripheral blood lymphocytes.
特記事項	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography. The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.4 Preservative: 0.097% Sodium azide Constituents: PBS, 0.2% BSA
精製度	Size exclusion

ポリモノ	モノクローナル
クローン名	B2M-01
アイソタイプ	IgG2a

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab49424の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration.

ターゲット情報

機能	Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system.
関連疾患	Defects in B2M are the cause of hypercatabolic hypoproteinemia (HYCATHYP) [MIM:241600]. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation. Note=Beta-2-microglobulin may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. Persistently high beta(2)-microglobulin serum levels lead to amyloidosis in patients on long-term hemodialysis.
配列類似性	Belongs to the beta-2-microglobulin family. Contains 1 Ig-like C1-type (immunoglobulin-like) domain.
翻訳後修飾	Glycation of Ile-21 is observed in long-term hemodialysis patients.
細胞内局在	Secreted. Detected in serum and urine.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors