

PE Anti-CD1a antibody [NA1/34] ab22352

2 References

製品の概要

製品名	PE Anti-CD1a antibody [NA1/34]
製品の詳細	PE Mouse monoclonal [NA1/34] to CD1a
由来種	Mouse
標識	PE. Ex: 488nm, Em: 575nm
特異性	ab222352 recognises specifically CD1a expressed by human cortical thymocytes, Langerhans cells, Dendritic cells and a subset of B cells.
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Human, Cynomolgus monkey
免疫原	Tissue, cells or virus corresponding to Human CD1a. Fresh human thymocytes
特記事項	<p>Recommended negative control: Mouse IgG2a Negative Control</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.20 Preservative: 0.09% Sodium azide Constituents: PBS, 1% BSA
特記事項(精製)	Purified IgG prepared by ion exchange chromatography
ポリ/モノ	モノクローナル
クローン名	NA1/34
ミエローマ	NS1/1-Ag4-1
アイソタイプ	IgG2a

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab22352の使用に適用されず
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 10µl for 10 ⁶ cells. ab91363 - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody. Use 10µl for 10 ⁶ cells in 100µl.

ターゲット情報

機能	Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.
組織特異性	Expressed on cortical thymocytes, epidermal Langerhans cells, dendritic cells, on certain T-cell leukemias, and in various other tissues.
配列類似性	Contains 1 Ig-like (immunoglobulin-like) domain.
細胞内局在	Cell membrane. Endosome membrane. Subject to intracellular trafficking between the cell membrane and endosomes. Localizes to cell surface lipid rafts.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors