

PE Anti-CD200R antibody [OX-110] ab35327

画像数 1

製品の概要

製品名	PE Anti-CD200R antibody [OX-110]
製品の詳細	PE Rat monoclonal [OX-110] to CD200R
由来種	Rat
標識	PE. Ex: 488nm, Em: 575nm
特異性	ab35327 recognises mouse CD200R, a cell surface glycoprotein (also known as OX2R).
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Mouse
免疫原	Fusion protein corresponding to CD200R. Database link: Q9ES57
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.40 Preservative: 0.09% Sodium azide Constituents: 1% BSA, PBS
精製度	Protein G purified
特記事項(精製)	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant.
ポリ/モノ	モノクローナル
クローン名	OX-110
アイソタイプ	IgG2a

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab35327の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		1/1. Use 10µl of this concentration to label 10 ⁶ cells in 100µl. ab134674 - Rat monoclonal IgG2a is suitable for use as an

ターゲット情報

機能	Receptor for the CD200/OX2 cell surface glycoprotein. Also binds to HHV-8 K14 viral CD200 homolog with identical affinity and kinetics as the host CD200.
組織特異性	Expressed in granulocytes, monocytes, most T cells, neutrophils, basophils and a subset of NK, NKT and B cells (at protein level). Expressed in bone marrow, lymph nodes, spleen, lung, liver, spinal cord, kidney. Expressed in monocyte-derived dendritic and mast cells.
配列類似性	Belongs to the CD200R family. Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
翻訳後修飾	The mature form of isoform 2 and/or isoform 4 starts at sequence position 27 of the corresponding isoform.
細胞内局在	Secreted and Cell membrane.

画像



ab35327 staining CD200R in mouse peripheral blood granulocytes by Flow Cytometric analysis.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors