abcam

Product datasheet

PE Anti-COX1 / Cyclooxygenase 1 antibody [EPR5866] ab209581

יובעבלא RabMAb

画像数 2

製品の概要

製品名 PE Anti-COX1 / Cyclooxygenase 1 antibody [EPR5866]

製品の詳細 PE Rabbit monoclonal [EPR5866] to COX1 / Cyclooxygenase 1

由来種 Rabbit

標識 PE. Ex: 488nm, Em: 575nm

アプリケーション 適用あり: Flow Cyt (Intra)

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール Flow Cyt (intra): HeLa cells

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at 4°C (stable for up to 12 months). Upon delivery aliquot. Store at +4°C.

Do Not Freeze. Store In the Dark.

バッファー

Preservative: 0.02% Sodium azide Constituents: 1% BSA, PBS

Protein A purified 精製度

ポリモノ モノクローナル

アイソタイプ

lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab209581の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/500.

ターゲット情報

機能	May play an important role in regulating or promoting cell proliferation in some normal and neoplastically transformed cells.	
パスウェイ	Lipid metabolism; prostaglandin biosynthesis.	
配列類似性	Belongs to the prostaglandin G/H synthase family.	

細胞内局在 Microsome membrane. Endoplasmic reticulum membrane.


画像



Flow Cytometry (Intracellular) - PE Anti-COX1 / Cyclooxygenase 1 antibody [EPR5866] (ab209581) Overlay histogram showing HeLa cells stained with ab209581 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilized with 90% methanol for 30 min at -20°C. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab209581, 1/500 dilution) for 30 min at 22°C.

Isotype control antibody (black line) was rabbit IgG (monoclonal) Phycoerythrin (<u>ab209478</u>) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 585/40 bandpass filter.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors