abcam

Product datasheet

PE/Cy7® Anti-CD45RA antibody [MEM-56] ab239312

画像数1

製品の概要

製品名 PE/Cy7® Anti-CD45RA antibody [MEM-56]

製品の詳細 PE/Cy7® Mouse monoclonal [MEM-56] to CD45RA

由来種 Mouse

標識 PE/Cy7®. Ex: 496nm, Em: 774nm

アプリケーション 適用あり: Flow Cyt

種交差性 交差種: Human

免疫原 Tissue, cells or virus corresponding to Human CD45RA. Human thymocytes and T lymphocytes.

Database link: P08575-8

ポジティブ・コントロール Flow Cyt: Human peripheral blood cells.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

クローン名

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Store In the Dark.

MEM-56

バッファー pH: 7.4

Preservative: 0.0975% Sodium azide

Constituent: PBS

精製度 Size exclusion

特記事項(精製) The purified antibody is conjugated with tandem dye PE-Cy™7 under optimum conditions. The

conjugate is purified by size-exclusion chromatography and adjusted for direct use. No

reconstitution is necessary.

ポリ/モノ モノクローナル

1

アプリケーション


The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab239312の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 4µl for 10 ⁶ cells. (or 100 µl of whole blood)

ターゲット情報

機能	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity.	
関連疾患	Severe combined immunodeficiency autosomal recessive T-cell-negative/B-cell-positive/NK-cell-positive Multiple sclerosis	
配列類似性	Belongs to the protein-tyrosine phosphatase family. Receptor class 1/6 subfamily. Contains 2 fibronectin type-Ill domains. Contains 2 tyrosine-protein phosphatase domains.	
ドメイン	The first PTPase domain interacts with SKAP1.	
翻訳後修飾	Heavily N- and O-glycosylated.	
細胞内局在	Membrane. Membrane raft. Colocalized with DPP4 in membrane rafts.	

画像



Flow cytometric analysis of human peripheral blood cells labeling CD45RA using ab239312. Surface staining. Gated on leukocytes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors