

### PerCP/Cy5.5® Anti-CD163 antibody [GHI/61] ab239338

#### 画像数 1

#### 製品の概要

---

|              |   |
|--------------|---|
| 製品名          | PerCP/Cy5.5® Anti-CD163 antibody [GHI/61]   |
| 製品の詳細        | PerCP/Cy5.5® Mouse monoclonal [GHI/61] to CD163   |
| 由来種          | Mouse   |
| 標識           | PerCP/Cy5.5®. Ex: 482nm, Em: 690nm  |
| アプリケーション     | <b>適用あり:</b> Flow Cyt   |
| 種交差性         | <b>交差種:</b> Human   |
| 免疫原          | Tissue, cells or virus corresponding to Human CD163. Hairy cell leukemia cells.   |
| ポジティブ・コントロール | Flow Cyt: Human peripheral blood.   |
| 特記事項         | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

#### 製品の特性

---

|          |  |
|----------|--|
| 製品の状態    | Liquid   |
| 保存方法     | Shipped at 4°C. Store at +4°C. Store In the Dark.  |
| バッファー    | pH: 7.4<br>Preservative: 0.0975% Sodium azide<br>Constituent: PBS  |
| 精製度      | Size exclusion   |
| 特記事項(精製) | The purified antibody is conjugated with tandem dye PerCP-Cy™5.5 under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary. |
| ポリ/モノ    | モノクローナル  |
| クローン名    | GHI/61   |
| アイソタイプ   | IgG1   |

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab239338の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項   |
|----------|-----------|--|
| Flow Cyt |           | Use 4µl for 10 <sup>6</sup> cells. OR 100 µl of whole blood. |

## ターゲット情報

|              |  |
|--------------|--|
| <b>機能</b>    | Acute phase-regulated receptor involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH-dependent manner. Exhibits a higher affinity for complexes of hemoglobin and multimeric haptoglobin of HP*1F phenotype than for complexes of hemoglobin and dimeric haptoglobin of HP*1S phenotype. Induces a cascade of intracellular signals that involves tyrosine kinase-dependent calcium mobilization, inositol triphosphate production and secretion of IL6 and CSF1. Isoform 3 exhibits the higher capacity for ligand endocytosis and the more pronounced surface expression when expressed in cells. After shedding, the soluble form (sCD163) may play an anti-inflammatory role, and may be a valuable diagnostic parameter for monitoring macrophage activation in inflammatory conditions. |
| <b>組織特異性</b> | Expressed in monocytes and mature macrophages such as Kupffer cells in the liver, red pulp macrophages in the spleen, cortical macrophages in the thymus, resident bone marrow macrophages and meningeal macrophages of the central nervous system. Expressed also in blood. Isoform 1 is the lowest abundant in the blood. Isoform 2 is the lowest abundant in the liver and the spleen. Isoform 3 is the predominant isoform detected in the blood.  |
| <b>配列類似性</b> | Contains 9 SRCR domains.   |
| <b>ドメイン</b>  | The SRCR domain 3 mediates calcium-sensitive interaction with hemoglobin/haptoglobin complexes.  |
| <b>翻訳後修飾</b> | A soluble form (sCD163) is produced by proteolytic shedding which can be induced by lipopolysaccharide, phorbol ester and Fc region of immunoglobulin gamma. This cleavage is dependent on protein kinase C and tyrosine kinases and can be blocked by protease inhibitors. The shedding is inhibited by the tissue inhibitor of metalloproteinase TIMP3, and thus probably induced by membrane-bound metalloproteinases ADAMs.<br>Phosphorylated.   |
| <b>細胞内局在</b> | Secreted and Cell membrane. Isoform 1 and isoform 2 show a lower surface expression when expressed in cells.   |

## 画像



Flow Cytometry analysis of human peripheral blood labeling CD163 with ab239338. Surface staining.

Flow Cytometry - PerCP/Cy5.5® Anti-CD163 antibody [GH1/61] (ab239338)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors