

Anti-Peripherin antibody [EPR23445-28] ab246502

リコンビナント RabMAb


★★★★★ **3 Abreviews** **1 References** **画像数 14**

製品の概要

製品名	Anti-Peripherin antibody [EPR23445-28]
製品の詳細	Rabbit monoclonal [EPR23445-28] to Peripherin
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IP, IHC-P, IHC-Fr, ICC/IF
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human spinal cord, Human lower limb nerve, Human nerve, SH-SY5Y, Mouse dorsal ganglion, Rat dorsal ganglion, Mouse spinal cord, Rat spinal cord, Mouse sciatic nerve, Neuro-2a and PC-12 lysates. IHC-P: Human colon, Mouse colon and Rat colon tissues. IHC-Fr: Mouse and rat colon tissue. ICC: SH-SY5Y and Neuro-2a cells. Flow:: SH-SY5Y and Neuro-2a cells. IP: Human spinal cord lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル



Western blot - Anti-Peripherin antibody [EPR23445-28] (ab246502)

All lanes : Anti-Peripherin antibody [EPR23445-28] (ab246502) at 1/1000 dilution

Lane 1 : Mouse dorsal ganglion lysate

Lane 2 : Rat dorsal ganglion lysate

Lane 3 : Mouse spinal cord lysate

Lane 4 : Rat spinal cord lysate

Lane 5 : Mouse sciatic nerve lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 54 kDa

Observed band size: 28,45,56,58 kDa

Blocking and dilution buffer: 5% NFDm/TBST.

Exposure time: 26 seconds.

The 58KD, 56KD, 45KD and 28KD isoforms of peripherin observed are consistent with what has been described in the literature (PMID: 20587592,18287500,17475883).



Western blot - Anti-Peripherin antibody [EPR23445-28] (ab246502)

All lanes : Anti-Peripherin antibody [EPR23445-28] (ab246502) at 1/1000 dilution

Lane 1 : Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Lane 2 : PC-12 (rat adrenal gland pheochromocytoma) whole cell lysate at 10 µg

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Predicted band size: 54 kDa

Observed band size: 45,56,58 kDa

Blocking and dilution buffer: 5% NFDm/TBST.

Exposure time: 6 seconds.

The 58KD, 56KD and 45KD isoforms of peripherin observed are consistent with what has been described in the literature (PMID: 20587592, 18287500, 17475883).



Western blot - Anti-Peripherin antibody [EPR23445-28] (ab246502)

All lanes : Anti-Peripherin antibody [EPR23445-28] (ab246502) at 1/1000 dilution

Lane 1 : Human spinal cord lysate

Lane 2 : Human lower limb nerve lysate

Lane 3 : Human nerve lysate

Lane 4 : SH-SY5Y (human neuroblastoma epithelial cell) whole cell lysate

Lane 5 : Human kidney lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) or VeriBlot for IP secondary antibody (HRP) ([ab131366](#)) at 1/100000 dilution

Predicted band size: 54 kDa


Observed band size: 45,56,58 kDa

Blocking and dilution buffer: 5% NFDm/TBST.

Exposure time: 37 seconds.

The 58KD, 56KD, and 45KD isoforms of peripherin observed are consistent with what has been described in the literature (PMID: 20587592, 18287500, 17475883).

Negative control: Kidney (PMID: 15132161).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Immunohistochemical analysis of paraffin-embedded Rat colon tissue labeling Peripherin with ab246502 at 1/10,000 dilution (0.05 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on the ganglion cells in rat colon. The section was incubated with ab246502 for 10 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND[®] RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling Peripherin with ab246502 at 1/10,000 dilution (0.05 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on the ganglion cells in mouse colon. The section was incubated with ab246502 for 10 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND[®] RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling Peripherin with ab246502 at 1/2000 dilution (0.227 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**). Positive staining on the ganglion cells in human colon (PMID: 26469323). The section was incubated with ab246502 for 10 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Immunoprecipitation - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Peripherin was immunoprecipitated from 0.35 mg Human spinal cord lysate with ab246502 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab246502 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used at 1/5000 dilution.

Lane 1: Human spinal cord lysate 10ug


Lane 2: ab246502 IP in Human spinal cord lysate

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab246502 in Human spinal cord lysate

Blocking and dilution buffer and concentration: 5% NFDN/TBST.


Exposure time: 40 seconds.

The 58KD isoform of peripherin observed is consistent with what has been described in the literature (PMID: 20587592, 18287500, 17475883).




Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized Neuro-2a (mouse neuroblastoma neuroblast) cells labelling Peripherin with ab246502 at 1/50 dilution (Red) compared with a Rabbit monoclonal IgG (**ab172730**) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor[®] 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.

Flow Cytometry (Intracellular) - Anti-Peripherin antibody [EPR23445-28] (ab246502)



Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized SH-SY5Y (human neuroblastoma epithelial cell) cells labelling Peripherin with ab246502 at 1/50 dilution (Red) compared with a Rabbit monoclonal IgG (**ab172730**) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor[®] 488, **ab150077**) at 1/2000 dilution was used as the secondary antibody.

Flow Cytometry (Intracellular) - Anti-Peripherin antibody [EPR23445-28] (ab246502)




Immunohistochemistry (Frozen sections) - Anti-
 Peripherin antibody [EPR23445-28] (ab246502)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized rat colon tissue labeling Peripherin with ab246502 at 1/250 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Positive staining on the ganglion cells in rat colon is observed. The nuclear counter stain is DAPI (Blue).

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.




Immunohistochemistry (Frozen sections) - Anti-
 Peripherin antibody [EPR23445-28] (ab246502)

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized mouse colon tissue labeling Peripherin with ab246502 at 1/250 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Positive staining on the ganglion cells in mouse colon is observed. The nuclear counter stain is DAPI (Blue).

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).


Secondary antibody only control: Used PBS instead of primary antibody, secondary is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Neuro-2a (mouse neuroblastoma neuroblast) cells labeling Peripherin with ab246502 at 1/100 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in Neuro-2a cell line. Tubulin is detected using **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution (Red). The nuclear counter stain is DAPI (Blue).

Secondary antibody only control: Used PBS instead of primary antibody, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Peripherin antibody [EPR23445-28] (ab246502)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized SH-SY5Y (human neuroblastoma epithelial cell) cells labeling Peripherin with ab246502 at 1/100 dilution, followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in SH-SY5Y cell line. Tubulin is detected using **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution (Red). The nuclear counter stain is DAPI (Blue).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit at 1/1000 dilution.

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-Peripherin antibody [EPR23445-28] (ab246502)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors