


### Anti-PHD3 antibody ab30782

★★★★★ **3 Abreviews** **17 References** 画像数 3

#### 製品の概要

製品名	Anti-PHD3 antibody
製品の詳細	Rabbit polyclonal to PHD3
由来種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, WB
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat 
免疫原	Synthetic peptide corresponding to Human PHD3 aa 50-150 (C terminal) conjugated to keyhole limpet haemocyanin. (Peptide available as <b>ab30781</b> )
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS  Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル

## アプリケーション

### The Abpromise guarantee

**Abpromise保証は、**次のテスト済みアプリケーションにおけるab30782の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 27 kDa (predicted molecular weight: 27 kDa).

## ターゲット情報

### 機能

Catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates HIF-1 alpha at 'Pro-564', and HIF-2 alpha. Functions as a cellular oxygen sensor and, under normoxic conditions, targets HIF through the hydroxylation for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. May play a role in cell growth regulation in muscle cells and in apoptosis in neuronal tissue. Promotes cell death through a caspase-dependent mechanism.

### 組織特異性

Widely expressed at low levels. Expressed at higher levels in heart (cardiac myocytes, aortic endothelial cells and coronary artery smooth muscle) and placenta.


### 配列類似性

Contains 1 Fe2OG dioxygenase domain.

### 細胞内局在

Cytoplasm. Nucleus.

## 画像



Western blot - Anti-PHD3 antibody (ab30782)

**All lanes :** Anti-PHD3 antibody (ab30782) at 1 µg/ml

**Lane 1 :** HeLa-Vehicle treated (Negative Control) Whole Cell Lysate ([ab116321](#))

**Lane 2 :** HeLa-DFO treated (0.5mM, 24h) Whole Cell Lysate ([ab116322](#))

**Lane 3 :** Human HeLa Nuclear DFO treated

**Lane 4 :** Human HeLa Cytoplasmic DFO treated

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


**Predicted band size:** 27 kDa

**Observed band size:** 27 kDa

**Additional bands at:** 18 kDa (possible non-specific binding)

**Exposure time:** 20 minutes

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 5% Bovine Serum Albumin before being incubated with ab30782 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution.



**All lanes :** Anti-PHD3 antibody (ab30782) at 1/1000 dilution

**Lane 1 :** Mouse embryonic fibroblast cell lysate. Treated 21% O<sub>2</sub> for 24 hours

**Lane 2 :** Mouse embryonic fibroblast cell lysate. Treated 1% O<sub>2</sub> for 24 hours

Lysates/proteins at 5 µg per lane.

#### **Secondary**

**All lanes :** Goat polyclonal anti-rabbit HRP-conjugate at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 27 kDa

**Exposure time:** 30 seconds



Immunocytochemistry/ Immunofluorescence - Anti-PHD3 antibody (ab30782)

ICC/IF image of ab30782 stained HepG2 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab30782 at 5µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti- rabbit (**ab96899**) IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors