abcam

Product datasheet

Anti-Phospholamban antibody [EPR21897] ab219626

ועלשעבע RabMAb

1 References 画像数7

製品の概要

製品名 Anti-Phospholamban antibody [EPR21897]

製品の詳細 Rabbit monoclonal [EPR21897] to Phospholamban

由来種 Rabbit

アプリケーション 適用あり: WB, IHC-P, IP

種交差性 交差種: Mouse, Rat, Human

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: Human heart and fetal heart tissue lysates; Mouse heart tissue lysate; Rat heart tissue lysate.

IHC-P: Human, mouse and rat cardiac muscle tissue. IP: Human fetal heart tissue lysate.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

精製度 Protein A purified

ポリモノ モノクローナル クローン名 EPR21897

アイソタイプ ΙgG


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab219626の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/2000. Detects a band of approximately 10, 25 kDa (predicted molecular weight: 6 kDa).
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/30.

ターゲット情報

機能	Reversibly inhibits the activity of ATP2A2 in cardiac sarcoplasmic reticulum by decreasing the apparent affinity of the ATPase for Ca(2+). Modulates the contractility of the heart muscle in response to physiological stimuli via its effects on ATP2A2. Modulates calcium re-uptake during muscle relaxation and plays an important role in calcium homeostasis in the heart muscle. The degree of ATP2A2 inhibition depends on the oligomeric state of PLN. ATP2A2 inhibition is alleviated by PLN phosphorylation.
組織特異性	Heart muscle (at protein level).
関連疾患	Cardiomyopathy, dilated 1P Cardiomyopathy, familial hypertrophic 18
配列類似性	Belongs to the phospholamban family.
翻訳後修飾	Phosphorylation by PKA abolishes the inhibition of ATP2A2-mediated calcium uptake. Phosphorylated at Thr-17 by CaMK2, and in response to beta-adrenergic stimulation. Phosphorylation by DMPK may stimulate sarcoplasmic reticulum calcium uptake in cardiomyocytes.
細胞内局在	Endoplasmic reticulum membrane. Sarcoplasmic reticulum membrane. Mitochondrion membrane. Membrane. Colocalizes with HAX1 at the endoplasmic reticulum (PubMed:17241641). Colocalizes with DMPK a the sarcoplasmic reticulum (PubMed:15598648).

画像



Western blot - Anti-Phospholamban antibody [EPR21897] (ab219626)

All lanes : Anti-Phospholamban antibody [EPR21897] (ab219626) at 1/2000 dilution

Lane 1: Human heart tissue lysate

Lane 2: Human fetal heart tissue lysate

Lane 3 : Mouse heart tissue lysate Lanes 4-5 : Rat heart tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/100000 dilution

Developed using the ECL technique.

Predicted band size: 6 kDa


Observed band size: 10,25 kDa

Exposure time: 15 seconds

Blocking and dilution buffer: 5% NFDM/TBST.

WB bands are consistent with reference (PMID 20696159; PMID

22144574)




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Phospholamban antibody [EPR21897] (ab219626)

Immunohistochemical analysis of paraffin embedded rat cardiac muscle tissue labeling Phospholamban with ab219626 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on human cardiac muscle (PMID: 28759816, PMID: 27642167) is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Phospholamban antibody [EPR21897] (ab219626)

Negative control: No staining shown in human tonsil tissue.

Immunohistochemical analysis of paraffin embedded human tonsil tissue labeling Phospholamban with ab219626 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. No staining on human tonsil is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Phospholamban antibody [EPR21897] (ab219626)

Immunohistochemical analysis of paraffin embedded mouse cardiac muscle tissue labeling Phospholamban with ab219626 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on mouse cardiac muscle (PMID: 28759816, PMID: 27642167) is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).



Immunoprecipitation - Anti-Phospholamban antibody [EPR21897] (ab219626)


Phospholamban was immunoprecipitated from 0.35 mg of human fetal heart tissue lysate with ab219626 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab219626 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/1000 dilution.

Lane 1: Human fetal heart tissue lysate 10 µg (Input).

Lane 2: ab219626 IP in human fetal heart tissue lysate.

Lane 3: Rabbit monoclonal IgG (<u>ab172730</u>) instead of ab219626 in human fetal heart tissue lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST. Exposure time: 30 seconds.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Phospholamban antibody [EPR21897] (ab219626)

Immunohistochemical analysis of paraffin embedded human cardiac muscle tissue labeling Phospholamban with ab219626 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on human cardiac muscle (PMID: 28759816, PMID: 27642167) is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit lgG H&L (HRP) ready to use.

Heat mediated antigen retrieval using <u>ab93684</u> (Tris/EDTA buffer, pH 9.0).



(ab219626)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors