abcam

Product datasheet

Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y775) antibody ab41434

画像数1

製品の概要

製品名 Anti-Phospholipase C gamma 1/PLC-gamma-1 (phospho Y775) antibody

製品の詳細 Rabbit polyclonal to Phospholipase C gamma 1/PLC-gamma-1 (phospho Y775)

由来種 Rabbit

特異性 Phospho-Tyr775-\Phospholipase C gamma 1/PLC-gamma-1 specific. Immunogen sequence has

low homology to Phospholipase C gamma 2.

アプリケーション **適用あり**: WB

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide corresponding to Human Phospholipase C gamma 1/PLC-gamma-1 (phospho

Y775).

Database link: P19174

ポジティブ・コントロール Human Jurkat and A431 cells treated with pervanadate.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at 4°C (up to 6 months). Store at -20°C.

ארע"ע". Preservative: 0.05% Sodium azide

Constituents: PBS, 50% Glycerol, 0.1% BSA

精製度 Immunogen affinity purified

特記事項(精製) This antibody was cross-adsorbed to a non-specific phospho-tyrosine peptide, then affinity

purified using phospho-Phospholipase C gamma 1/PLC-gamma-1 (Tyr775) peptide.

ポリ/モノ ポリクローナル

アイソタイプ lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab41434の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 150 kDa (predicted molecular weight: 150 kDa).

ターゲット情報

機能 Plays a role in actin reorganization and cell migration. The production of the second messenger

molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. Major substrate for heparin-binding

growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase.

配列類似性 Contains 1 C2 domain.

Contains 1 EF-hand domain. Contains 2 PH domains.

Contains 1 PI-PLC X-box domain. Contains 1 PI-PLC Y-box domain.

Contains 2 SH2 domains. Contains 1 SH3 domain.

ドメイン The SH3 domain mediates interaction with CLNK (By similarity). The SH3 domain also mediates

interaction with RALGPS1.

翻訳後修飾 The receptor-mediated activation of PLC-gamma-1 and PLC-gamma-2 involves their

phosphorylation by tyrosine kinases in response to ligation of a variety of growth factor receptors


and immune system receptors. May be dephosphorylated by PTPRJ.

Ubiquitinated by CBLB in activated T-cells.

細胞内局在 Cell projection > lamellipodium. Cell projection > ruffle. Rapidly redistributed to ruffles and

lamellipodia structures in response to epidermal growth factor (EGF) treatment.

画像



Western blot - Anti-Phospholipase C gamma 1/PLCgamma-1 (phospho Y775) antibody (ab41434)

Western blot analysis of Phospholipase C gamma 1/PLC-gamma-1 immunoprecipitates from human jurkat cells untreated (lanes 1 & 3) or treated with pervanadate (1 mM) for 30 min (lanes 2, 4, 5 & 6). Immunoprecipitation was performed with monoclonal anti-Phospholipase C gamma 1/PLC-gamma-1. The blots were probed with anti-Phospholipase C gamma 1/PLC-gamma-1 (lanes 1 & 2) and anti-Phospholipase C gamma 1/PLC-gamma-1 (Tyr-775) (lanes 3-6). The latter antibody was used in the presence of phospho-Phospholipase C gamma 1/PLC-gamma-1 (Tyr-775) peptide (lane 5), or a non-specific phosphotyrosine peptide (lane 6).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors