

Anti-PKA R2/PKR2 (phospho S99) antibody [E151] - BSA and Azide free ab238951

リコンビナント **RabMAb**

画像数 5

製品の概要

製品名	Anti-PKA R2/PKR2 (phospho S99) antibody [E151] - BSA and Azide free
製品の詳細	Rabbit monoclonal [E151] to PKA R2/PKR2 (phospho S99) - BSA and Azide free
由来種	Rabbit
特異性	<p>The antibody only detects PKA R2/PKR2 phosphorylated on Serine 99. Phosphoserine 99 in Human corresponds to phosphoserine 96 in mouse.</p> <p>The mouse and pig recommendation is based on the WB results. We do not guarantee IHC-P for mouse and pig.</p>
アプリケーション	<p>適用あり: WB, ICC/IF, IHC-P</p> <p>適用なし: Flow Cyt</p>
種交差性	交差種: Mouse, Rat, Human, Pig
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	ICC/IF: HeLa cells. IHC-P: Human lymph node metastatic tissue. Human kidney without Lambda Protein Phosphatase treatment (image A) and Rat kidney without Lambda Protein Phosphatase treatment
特記事項	<p>ab238951 is the carrier-free version of ab32390.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	E151
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab238951の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Detects a band of approximately 51 kDa (predicted molecular weight: 45 kDa).
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. The mouse and pig recommendation is based on the WB results. We do not guarantee IHC-P for mouse and pig. See IHC antigen retrieval protocols .

追加情報 Is unsuitable for Flow Cyt.

ターゲット情報

機能	Regulatory subunit of the cAMP-dependent protein kinases involved in cAMP signaling in cells. Type II regulatory chains mediate membrane association by binding to anchoring proteins, including the MAP2 kinase.
組織特異性	Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression

配列類似性

varies among tissues and is in some cases constitutive and in others inducible.

翻訳後修飾

Belongs to the cAMP-dependent kinase regulatory chain family.


Contains 2 cyclic nucleotide-binding domains.

細胞内局在

Phosphorylated by the activated catalytic chain.


Cytoplasm. Cell membrane. Colocalizes with PJA2 in the cytoplasm and the cell membrane.

画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PKA R2/PKR2 (phospho S99) antibody [E151] - BSA and Azide free (ab238951)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat kidney without Lambda Protein Phosphatase treatment (image A) and treated with Lambda Protein Phosphatase (image B). tissue sections labeling PKA R2/PKR2 with purified [ab32390](#) at 1:500 dilution (0.75 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PKA R2/PKR2 (phospho S99) antibody [E151] - BSA and Azide free (ab238951)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human kidney without Lambda Protein Phosphatase treatment (image A) and treated with Lambda Protein Phosphatase (image B). tissue sections labeling PKA R2/PKR2 with purified [ab32390](#) at 1:500 dilution (0.75 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



ab32390, at a 1/250 dilution, staining (A) untreated (B) Phosphatase treated lymph node metastatic by immunohistochemistry, Paraffin embedded human tissue

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32390**).



Cell line: HeLa (human cervix adenocarcinoma)

Target AbID: Ab32390 anti-PKA R2/PKR2 (phospho S99)

Target Secondary: **ab150077** AlexaFluor®488 goat anti-rabbit

Counterstain AbID: **ab7291** anti-Tubulin (mouse mAb)

Target Secondary: **ab150120** AlexaFluor®594 goat anti-mouse

Fixative: 4% PFA

Permeabilization: 0.1% Triton-X

Nuclear counter stain: DAPI

Other comments: Confocal image showed increased cytoplasmic staining after CA (100ng/ml, 10min) treatment on Hela cells. The LP treatment decreased the increased cytoplasmic staining caused by CA. Ab32514 was used as a Pan control for **ab32390**. The results showed cytoplasmic staining on untreated, CA and CA+LP treated Hela cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32390**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-PKA R2/PKR2 (phospho S99) antibody [E151] -
BSA and Azide free (ab238951)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors