abcam

Product datasheet

Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free ab239844

יובעבלא RabMAb

画像数6

製品の概要

免疫原

特記事項

製品名 Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EP1511Y] to PPP1CA + PPP1CB - BSA and Azide free

由来種 Rabbit

特異性 The immunogen for this antibody is 100% homologous with Human PPP1CA and PPP1CB

アプリケーション 適用あり: IHC-P, Flow Cvt (Intra), IP, ICC/IF, WB

種交差性 交差種: Mouse, Rat, Human

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ab239844 is the carrier-free version of ab52619.

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

バッファー pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

ポリ/モノ モノクローナル **クローン名** EP1511Y

アイソタイプ IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab239844の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols.
Flow Cyt (Intra)		Use at an assay dependent concentration. <u>ab199376</u> - Rabbit monoclonal lgG, is suitable for use as an isotype control with this antibody.
IP		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Predicted molecular weight: 37 kDa.

ターゲット情報

細胞内局在 Cytoplasm. Nucleus. Nucleus > nucleoplasm. Nucleus > nucleolus.


画像



Immunocytochemistry/ Immunofluorescence - Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free (ab239844)

ab52619 staining PPP1CA + 1CB in the HepG2 cell line by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with primary antibody (1/50). **ab150077**(1/500) an Alexa Fluor[®]488-conjugated Goat anti-rabbit IgG was used as the secondary antibody. Nuclei were counterstained with DAPI.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab52619).




Immunoprecipitation - Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free (ab239844)

<u>ab52619</u> (purified) at 1/30 immunoprecipitating PPP1CA + 1CB in Jurkat cell lysate. For western blotting, a HRP-conjugated Goat antirabbit lgG, specific to the non-reduced form of lgG was used as the secondary antibody (1/1000).

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab52619</u>).



Flow Cytometry (Intracellular) - Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free (ab239844)

Overlay histogram showing HeLa cells stained with <u>ab52619</u> (red line) at 1/150 dilution. The cells were fixed with 80% methanol. The secondary antibody used was a FITC conjugated goat anti-rabbit IgG at 1/150 dilution. Isotype control antibody (black line) was rabbit monoclonal IgG used under the same conditions. Cells also incubated without primary antibody and secondary antibody (blue line)


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab52619).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free (ab239844)

<u>ab52619</u> staining PP1CA + 1CB in Human cerebrum cortex tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed and paraffinembedded, antigen retrieval was by heat mediation in Tris/EDTA buffer pH9. Samples were incubated with primary antibody (1/50). An undiluted HRP-conjugated mouse anti-rabbit lgG was used as the secondary antibody. Tissue counterstained with Hematoxylin. PBS was used in the negative control rather than the Primary antibody.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab52619</u>).



Flow Cytometry (Intracellular) - Anti-PPP1CA + PPP1CB antibody [EP1511Y] - BSA and Azide free (ab239844)

Overlay histogram showing HeLa cells stained with <u>ab52619</u>, unpurified (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (<u>ab52619</u>, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight[®] 488 goat antirabbit lgG (H+L) (<u>ab96899</u>) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit lgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab52619).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors