

### Anti-pro Caspase-3 antibody [E83-103] ab32499

KO 評価済 リコンビナント RabMAb

79 References 画像数 6

#### 製品の概要

製品名	Anti-pro Caspase-3 antibody [E83-103]
製品の詳細	Rabbit monoclonal [E83-103] to pro Caspase-3
由来種	Rabbit
特異性	This antibody only detects pro-form (35kD) of caspase-3, and does not recognize any cleaved caspases.
アプリケーション	<b>適用あり:</b> ICC/IF, WB, IHC-P, Flow Cyt (Intra)
種交差性	<b>交差種:</b> Mouse, Human
免疫原	Synthetic peptide corresponding to Human pro Caspase-3 (N terminal).
ポジティブ・コントロール	Jurkat whole cell lysate ( <b>ab7899</b> ) and human colon adenocarcinoma.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <b><a href="#">see here</a></b>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <b><a href="#">RabMAb<sup>®</sup> patents</a></b>.</p> <p>Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル

クローン名 E83-103  
アイソタイプ IgG

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab32499の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
WB		1/10000. Detects a band of approximately 35 kDa (predicted molecular weight: 31 kDa).
IHC-P		1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Flow Cyt (Intra)		1/50. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.


## ターゲット情報

**関連性**

Caspases are a family of cysteine proteases that are key mediators of programmed cell death or apoptosis. The precursor form of all caspases is composed of a prodomain, and large and small catalytic subunits. The active forms of caspases are generated by several stimuli including ligand-receptor interactions, growth factor deprivation and inhibitors of cellular functions. All known caspases require cleavage adjacent to aspartates to liberate one large and one small subunit, which associate into a2b2 tetramer to form the active enzyme. Gene for Caspase 3 also known as Yama, CPP32, and apopain codes for a 32-kDa protein. Caspase 3 cleaves the death substrate poly(ADP-ribose) polymerase (PARP) to a specific 85 kDa form observed during apoptosis and is inhibitable by the CrmA protein. Other Caspase 3 substrates include DNA-PK, actin, GAS2, and procaspase-6, etc. Caspase 3 is activated by cleavage events at Asp-28/Ser-29 (between N-terminal pro-domain) and Asp-175/Ser-176 (between large and small subunits) to generate a large subunit of 17-kDa and a small subunit of 12-kDa.

**細胞内局在**      Cytoplasmic

## 画像



Western blot - Anti-pro Caspase-3 antibody [E83-103] (ab32499)

**Lane 1:** Wild-type HAP1 cell lysate


**Lane 2:** Wild-type HAP1 cell lysate + Staurosporine (1µM for 4h)

**Lane 3:** Caspase-3 knockout HAP1 cell lysate

**Lane 4:** Caspase-3 knockout HAP1 cell lysate + Staurosporine (1µM for 4h)

**Lanes 1 - 4:** Merged signal (red and green). Green - ab32499 observed at 35 kDa. Red - loading control, **ab8245**, observed at 37 kDa.


ab32499 was shown to specifically react with pro Caspase 3 when Caspase 3 knockout samples were used. Wild-type and Caspase 3 knockout samples ( $\pm$  Staurosporine treatment) were subjected to SDS-PAGE. ab32499 and **ab8245** (loading control to GAPDH) were diluted to 1/1000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-pro Caspase-3 antibody [E83-103] (ab32499)

Immunohistochemical analysis of paraffin-embedded human colon adenocarcinoma ab32499 at 1/250 dilution.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-pro Caspase-3 antibody [E83-103] (ab32499)



Flow Cytometry (Intracellular) - Anti-pro Caspase-3 antibody [E83-103] (ab32499)



Western blot - Anti-pro Caspase-3 antibody [E83-103] (ab32499)

ICC/IF image of ab32499 stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab32499, 5µg/ml) overnight at +4°C. The secondary antibody (green) was [ab96899](#), anti-rabbit DyLight® 488 used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Overlay histogram showing Jurkat cells stained with ab32499 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab32499, 1/50 dilution) for 30 min at 22°C. The secondary antibody used was anti-rabbit DyLight® 488 ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >5,000 events was performed.

**All lanes :** Anti-pro Caspase-3 antibody [E83-103] (ab32499) at 1/10000 dilution

**Lane 1 :** Jurkat cell lysate

**Lane 2 :** Jurkat cell lysate + Camptothecin

**Predicted band size:** 31 kDa

**Observed band size:** 35 kDa

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-pro Caspase-3 antibody [E83-103] (ab32499)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors