


Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker ab2723

★★★★★ [14 Abreviews](#) [169 References](#) [画像数 3](#)

製品の概要

製品名	Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker
製品の詳細	Mouse monoclonal [6G6-1C9] to PSD95 - Synaptic Marker
由来種	Mouse
特異性	We do not guarantee IHC-P for mouse.
アプリケーション	適用あり: IHC-P, WB
種交差性	交差種: Mouse, Rat 交差が予測される動物種: Cow, Zebrafish 
免疫原	Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Rat retina tissue. WB: Rat and mouse hippocampus and forebrain tissue lysate.
特記事項	<p>The Protocols tab contains a Mouse on Mouse staining protocol with recommendations when using a mouse monoclonal antibody to stain mouse tissues and tips for reducing background.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: PBS, 6.97% L-Arginine</p>
精製度	Protein G purified

ポリ/モノ	モノクローナル
クローン名	6G6-1C9
アイソタイプ	IgG2a

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab2723の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 1 - 2 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. We do not guarantee IHC-P for mouse.
WB	★★★★★ (7)	Use a concentration of 1 - 5 µg/ml. Detects a band of approximately 95 kDa (predicted molecular weight: 80 kDa).

ターゲット情報

機能	Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling. Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining the channel intracellularly. May regulate the intracellular trafficking of ADR1B.
組織特異性	Brain.
配列類似性	Belongs to the MAGUK family. Contains 1 guanylate kinase-like domain. Contains 3 PDZ (DHR) domains. Contains 1 SH3 domain.
ドメイン	The PDZ domain 3 mediates interaction with ADR1B. The L27 domain near the N-terminus of isoform 2 is required for HGS/HRS-dependent targeting to postsynaptic density.
翻訳後修飾	Palmitoylation of isoform 1 is required for targeting to postsynaptic density.
細胞内局在	Cell membrane. Cell junction, synapse, postsynaptic cell membrane, postsynaptic density. Cell projection, axon. Cell junction, synapse. High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on axon hillocks of Purkinje cells.

画像



Secondary-only control


Copyright (c) 2019 Abcam plc

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723)

IHC image of ab2723 staining in formalin-fixed, paraffin-embedded rat retinal tissue, performed on a Leica BOND™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab2723, 2µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX.

Secondary-only control image is shown as insert.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Western blot - Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723)

All lanes : Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723) at 5 µg/ml

Lane 1 : Hippocampus (Rat) Tissue Lysate

Lane 2 : Forebrain (Rat) Tissue Lysate

Lane 3 : Hippocampus (Mouse) Tissue Lysate

Lane 4 : Forebrain (Mouse) Tissue Lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/5000 dilution


Performed under reducing conditions.

Predicted band size: 80 kDa

Observed band size: 95 kDa

Additional bands at: 100 kDa, 110 kDa, 80 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 3 minutes



Western blot - Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723)

Image courtesy of an AbReview submitted by Grant Corbett

Lanes 1-6 : Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723) at 1/1200 dilution

Lanes 7-12 : Anti-PSD95 antibody [6G6-1C9] - Synaptic Marker (ab2723) at 1/1200 dilution (18 hours at 4°C)

Lanes 1-6 : Mouse Hippocampus with Li-Cor Block Buffer, 1 hour at 21°C

Lanes 7-12 : Mouse Liver with Li-Cor Block Buffer, 1 hour at 21°C

Lysates/proteins at 30 µg per lane.

Secondary

All lanes : Li-Cor IRDye® Donkey anti-mouse 680LT at 1/18000 dilution

Performed under reducing conditions.

Predicted band size: 80 kDa

Additional bands at: 95 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 5 minutes

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors