


Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1] ab241062

リコンビナント RabMAb

5 References 画像数 5

製品の概要

製品名	Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1]
製品の詳細	Rabbit monoclonal [MJF-R24-17-1] to RAB29 (phospho T71)
由来種	Rabbit
アプリケーション	適用あり: Dot blot, WB
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C] whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>This antibody was developed with support from The Michael J. Fox Foundation.</p>



製品の特性

製品の状態	Liquid
-------	--------

保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリモノ	モノクローナル
クローン名	MJF-R24-17-1
アイソタイプ	IgG


アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab241062の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Dot blot		1/1000.
WB		1/500. Detects a band of approximately 23 kDa (predicted molecular weight: 23 kDa).

ターゲット情報

機能	Rab GTPase key regulator in vesicle trafficking. Essential for maintaining the integrity of the endosome-trans-Golgi network structure. Together with LRRK2, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner. Regulates neuronal process morphology in the intact central nervous system (CNS). May play a role in the formation of typhoid toxin transport intermediates during Salmonella enterica serovar Typhi (S.Typhi) epithelial cell infection.
組織特異性	Ubiquitous.
配列類似性	Belongs to the small GTPase superfamily. Rab family.
翻訳後修飾	In case of Salmonella enterica serovar Typhimurium (S.Typhimurium) infection, is proteolytically cleaved between Gly-41 and Val-42 by the GtgE viral protease encoded on the Gifsy-2 lysogen bacteriophage, which therefore prevents the recruitment of RAB29 to S.Typhimurium-containing vacuoles. In contrast, no proteolytically cleavage is detected in S.Typhi-infected cells (PubMed:22042847).
細胞内局在	Cell membrane. Cytoplasm. Cytoplasm, perinuclear region. Golgi apparatus. Golgi apparatus, trans-Golgi network. Vacuole. Cytoplasm, cytoskeleton. Colocalizes with LRRK2 along tubular structures emerging from Golgi apparatus (By similarity). Colocalizes with GM130 at the Golgi apparatus. Colocalizes with dynamic tubules emerging from and retracting to the Golgi apparatus. Colocalizes with TGN46 at the trans-Golgi network (TGN). In Salmonella enterica serovar Typhi (S.Typhi) infected epithelial cells, is recruited and colocalized with both S.Typhi-containing vacuoles and dynamic tubules as well as those emerging from the vacuole toward the cell periphery.



All lanes : Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1] (ab241062) at 1/5000 dilution

Lane 1 : HEK-293 (human embryonic kidney) over-expressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate

Lane 2 : MLi-2 (100nM for 60min) treated HEK-293 over-expressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate

Lysates/proteins at 15 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 23 kDa

Observed band size: 23 kDa

Blocking buffer: 5% NFDM/TBST.



Dot blot analysis of RAB29 (phospho T71) labeled with ab241062 at 1/1000 dilution.


Lane 1: RAB29 (phospho T71) peptide.

Lane 2: RAB29 non-phospho peptide.

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution was used as secondary antibody.

Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.



All lanes : Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1] (ab241062) at 1/500 dilution

Lane 1 : HEK-293 (human epithelial cell line from embryonic kidney) overexpressing HA-tagged RAB3 and LRRK2 [Y1699C], whole cell lysate

Lane 2 : HEK-293 overexpressing HA-tagged RAB5 and LRRK2 [Y1699C], whole cell lysate

Lane 3 : HEK-293 overexpressing HA-tagged RAB8 and LRRK2 [Y1699C], whole cell lysate

Lane 4 : HEK-293 overexpressing HA-tagged RAB10 and LRRK2 [Y1699C], whole cell lysate

Lane 5 : HEK-293 overexpressing HA-tagged RAB12 and LRRK2 [Y1699C], whole cell lysate

Lane 6 : HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate

Lane 7 : HEK-293 overexpressing HA-tagged RAB35 and LRRK2 [Y1699C], whole cell lysate

Lane 8 : HEK-293 overexpressing HA-tagged RAB43 and LRRK2 [Y1699C], whole cell lysate

Lysates/proteins at 0.5 µg per lane.

Secondary

All lanes : IRDye® 800CW Goat-anti-Rabbit at 1/25000 dilution

Predicted band size: 23 kDa


Observed band size: 23 kDa

Blocking buffer: 5% NFDM/TBST.

Dilution buffer: 5% BSA/TBST.

The blot was scanned with LICOR Odyssey CLx.

This image is kindly provided by our collaborator Dr. Dario Alessi, University of Dundee.



Western blot - Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1] (ab241062)

All lanes : Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1] (ab241062) at 1/500 dilution

Lane 1 : MLI-2 (100nM for 60min) treated HEK-293 (human epithelial cell line from embryonic kidney) overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate at 3 µg

Lane 2 : HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate at 3 µg

Lane 3 : HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate at 1 µg

Lane 4 : HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate at 0.3 µg

Lane 5 : HEK-293 overexpressing HA-tagged RAB29 and LRRK2 [Y1699C], whole cell lysate at 0.1 µg

Secondary

All lanes : IRDye® 800CW Goat-anti-Rabbit at 1/25000 dilution

Predicted band size: 23 kDa

Observed band size: 23 kDa

Blocking buffer: 5% NFDM/TBST.

Dilution buffer: 5% BSA/TBST.

The LRRK2 pathogenic mutation Y1699C increases LRRK2 activity and markedly elevates the phosphorylation of Rab proteins. While MLI-2 is a potent, selective LARRK2 kinase inhibitor (PMID: 26407721).

The blot was scanned with LI-COR Odyssey CLx.

This image is kindly provided by our collaborator Dr. Dario Alessi, University of Dundee

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-RAB29 (phospho T71) antibody [MJF-R24-17-1]
(ab241062)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors