

Anti-Rab4 antibody [EPR3043] - Early Endosome Marker ab109009

KO 評価済 リコンビナント RabMAb

★★★★★ [1 Abreviews](#) [10 References](#) [画像数 9](#)

製品の概要

製品名	Anti-Rab4 antibody [EPR3043] - Early Endosome Marker
製品の詳細	Rabbit monoclonal [EPR3043] to Rab4 - Early Endosome Marker
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IP, ICC/IF 適用なし: IHC-P
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: MCF7, PC12, Neuro 2a, 293T, SH SY5Y and Human fetal brain lysates; ICC/IF: HeLa cells. Flow Cyt (intra): HeLa cells. IP: MCF7 cells.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), PBS, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR3043

アイソタイプ

IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab109009の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/200.
WB		1/1000 - 1/10000. Predicted molecular weight: 24 kDa.
IP		1/10 - 1/100.
ICC/IF	★☆☆☆☆ (1)	1/170 - 1/1000.


追加情報

Is unsuitable for IHC-P.

ターゲット情報


機能	Protein transport. Probably involved in vesicular traffic.
配列類似性	Belongs to the small GTPase superfamily. Rab family.
翻訳後修飾	Phosphorylated by CDK1 kinase during mitosis.
細胞内局在	Membrane. Cytoplasm. Generally associated with membranes. Cytoplasmic when phosphorylated by CDK1.

画像



Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Rab4 with Purified ab109009 at 1:170 dilution (10 µg/ml). Cells were fixed in 100% Methanol. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

Immunocytochemistry/ Immunofluorescence - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)



Western blot - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: Rab4 knockout HAP1 cell lysate (20 µg)

Lane 3: HeLa cell lysate (20 µg)

Lane 4: Human fetal brain lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab109009 observed at 24 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab109009 was shown to specifically react with Rab4 when Rab4 knockout samples were used. Wild-type and Rab4 knockout samples were subjected to SDS-PAGE. ab109009 and **ab8245** (loading control to GAPDH) were diluted 1/1000 and 1/2000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/1000 dilution (Purified)

Lane 1 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates

Lane 2 : Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysates


Lane 3 : MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

Lysates/proteins at 1/15 dilution per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 24 kDa



Immunoprecipitation - Anti-Rab4 antibody
 [EPR3043] - Early Endosome Marker (ab109009)

ab109009 (purified) at 1:80 dilution (2µg) immunoprecipitating Rab4 in MCF7 whole cell lysate.


Lane 1 (input): MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate 10µg

Lane 2 (+): ab109009 & MCF7 whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab109009 in MCF7 whole cell lysate


For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.




Flow Cytometry (Intracellular) - Anti-Rab4 antibody
 [EPR3043] - Early Endosome Marker (ab109009)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Rab4 with Purified ab109009 at 1/200 dilution (1µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunocytochemistry/ Immunofluorescence - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

Immunocytochemistry/Immunofluorescence analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) labelling Rab4 with purified ab109009 at 1/250. Cells were fixed with 100% methanol and permeabilized with 0.1% triton X-100. **ab150077** Goat anti rabbit IgG (Alexa Fluor® 488) at 1/1000 was used as the secondary antibody. Nuclei were counterstained with DAPI. PBS was used instead of the primary antibody as the negative control.



Western blot - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

All lanes : Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009) at 1/1000 dilution

Lane 1 : MCF7 cell lysate

Lane 2 : PC12 cell lysate

Lane 3 : Neuro 2a cell lysate


Lane 4 : 293T cell lysate

Lane 5 : SH SY5Y cell lysate

Lane 6 : Human fetal brain lysate

Lysates/proteins at 10 µg per lane.


Predicted band size: 24 kDa



ab109009 at 1/500 dilution staining Rab4 in HeLa by Immunofluorescence.

Immunocytochemistry/ Immunofluorescence - Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-Rab4 antibody [EPR3043] - Early Endosome Marker (ab109009)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors