


Anti-RUNX2 antibody [EPR14334] ab192256

リコンビナント RabMAb

★★★★★ **8 Abreviews** **92 References** 画像数 11

製品の概要

製品名	Anti-RUNX2 antibody [EPR14334]
製品の詳細	Rabbit monoclonal [EPR14334] to RUNX2
由来種	Rabbit
アプリケーション	適用あり: IHC-P, ICC/IF, Flow Cyt (Intra), ChIC/CUT&RUN-seq
種交差性	交差種: Mouse, Human 交差が予測される動物種: Rat 
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Human osteosarcoma, Human tonsil and Mouse spleen tissues; Saos-2 and PC cells.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
精製度	Protein A purified
ポリモノ	モノクローナル
クローン名	EPR14334
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab192256の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (5)	1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/1000. For unpurified use at 1/500.
Flow Cyt (Intra)		1/50.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration.

ターゲット情報

機能	Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis. Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'-PYGPGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters (By similarity). Inhibits MYST4-dependent transcriptional activation.
組織特異性	Specifically expressed in osteoblasts.
関連疾患	Defects in RUNX2 are the cause of cleidocranial dysplasia (CLCD) [MIM:119600]; also known as cleidocranial dysostosis (CCD). CLCD is an autosomal dominant skeletal disorder with high penetrance and variable expressivity. It is due to defective endochondral and intramembranous bone formation. Typical features include hypoplasia/aplasia of clavicles, patent fontanelles, wormian bones (additional cranial plates caused by abnormal ossification of the calvaria), supernumerary teeth, short stature, and other skeletal changes. In some cases defects in RUNX2 are exclusively associated with dental anomalies.
配列類似性	Contains 1 Runt domain.
ドメイン	A proline/serine/threonine rich region at the C-terminus is necessary for transcriptional activation of target genes and contains the phosphorylation sites.
翻訳後修飾	Phosphorylated; probably by MAP kinases (MAPK) (By similarity). Isoform 3 is phosphorylated on Ser-340.
細胞内局在	Nucleus.

画像



Western blot - Anti-RUNX2 antibody [EPR14334] (ab192256)

All lanes : Anti-RUNX2 antibody [EPR14334] (ab192256) at 1/1000 dilution

Lane 1 : Saos-2 cell lysate

Lane 2 : MC3T3-E1 undifferentiated cell lysate

Lane 3 : MC3T3-E1 7-day Osteogenic differentiation cell lysate

Lane 4 : MC3T3-E1 14-day Osteogenic differentiation cell lysate

Lane 5 : MC3T3-E1 28-day Osteogenic differentiation cell lysate

Lane 6 : C2C12 cell lysate

Lane 7 : SH-SY5Y cell lysate


Lane 8 : NIH/3T3 cell lysate

Lane 9 : LNCaP cell lysate

Lysates/proteins at 20 µg per lane.

Observed band size: 60 kDa


Western blot: Anti-RUNX2 antibody [EPR14334] (ab192256) staining at 1/1000 dilution, shown in green; Mouse anti-CANX [CANX/1543] (**ab238078**) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab192256 was shown to bind specifically to RUNX2. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution



ChIP/CUT&RUN sequencing - Anti-RUNX2 antibody [EPR14334] (ab192256)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/μL, 2.5 x 10⁵ Saos-2 cells and 5 μg of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.



ChIP/CUT&RUN sequencing - Anti-RUNX2 antibody [EPR14334] (ab192256)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/μL, 2.5 x 10⁵ Saos-2 cells and 5 μg of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.



ChIP/CUT&RUN sequencing - Anti-RUNX2 antibody [EPR14334] (ab192256)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/ μ L, 2.5×10^5 Saos-2 cells and 5 μ g of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody [EPR14334] (ab192256)


Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-RUNX2 antibody [EPR14334] (ab192256)


Immunocytochemistry/Immunofluorescence analysis of Saos-2 (Human osteosarcoma cell line) labeling RUNX2 with purified ab192256 at 1/1000 dilution. Cells were fixed with 4% PFA and permeabilized with 0.1% tritonX-100. **ab150077** Goat anti rabbit IgG (Alexa Fluor[®]488) at 1/1000 was used as the secondary antibody. Nuclei were counterstained with DAPI. PBS was used instead of the primary antibody as the negative control.



Immunohistochemical analysis of paraffin-embedded Human osteosarcoma tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody [EPR14334] (ab192256)




ab192256 staining RUNX2 in PC-3 (human prostate adenocarcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde, permeabilised with 90% methanol and the sample was incubated with the primary antibody at a dilution of 1/50. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isootype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)


Flow Cytometry (Intracellular) - Anti-RUNX2 antibody [EPR14334] (ab192256)



Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody [EPR14334] (ab192256)



Immunocytochemistry/ Immunofluorescence - Anti-RUNX2 antibody [EPR14334] (ab192256)

Immunofluorescent analysis of 4% formaldehyde fixed PC3 cells labeling RUNX2 using ab192256 at a 1/500 dilution. A Goat anti rabbit IgG (Alexa Fluor®488) **ab150077** was used as the secondary at a 1/200 dilution. Counterstain DAPI. Permeabilized using 0.1% Triton X-100. The two negative controls: 1. Primary ab concentration (anti-RUNX2) is 1:500 dilution, Secondary ab (Goat anti mouse IgG (Alexa Fluor®594)) is 1:500 dilution; 2. Primary ab concentration (anti-RUNX2) is 1:500 dilution, Secondary ab (Goat anti mouse IgG (Alexa Fluor®594)) is 1:500 dilution.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-RUNX2 antibody [EPR14334] (ab192256)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors