

Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal ab199719

KO 評価済 リコンビナント RabMAb

★★★★★ [1 Abreviews](#) [7 References](#) [画像数 8](#)

製品の概要

製品名	Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal
製品の詳細	Rabbit monoclonal [EPR17319] to SLC27A4 / FATP4 - C-terminal
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, IP, IHC-P, WB
種交差性	交差種: Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HeLa, HEK293 and HepG2 whole cell lysates; Human fetal brain and Human fetal kidney lysates. IHC-P: Human kidney tissue. ICC/IF: HepG2 cells. IP: HEK293 whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR17319

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab199719の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
ICC/IF		1/100.
IP		1/50.
IHC-P		1/50. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB	★★★★★ (1)	1/2000. Detects a band of approximately 72 kDa (predicted molecular weight: 72 kDa).

ターゲット情報

関連性 SLC27A4 / FATP4 plays a role in the transport of long chain fatty acids across the plasma membrane. It has acyl-coA ligase activity for long chain and very long chain fatty acids. Deletion of the SLC27A4 gene results in embryonic lethality, which is attributed to the need for fat absorption across the visceral endoderm early in embryonic development. Expression of FAT4P in the intestinal lining is thought to be altered in response to dietary fat.

細胞内局在 Cell Membrane

画像



All lanes : Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719) at 1/1000 dilution

- Lane 1 :** Wild-type HEK293T cell lysate
- Lane 2 :** SLC27A4 knockout HEK293T cell lysate
- Lane 3 :** HepG2 cell lysate
- Lane 4 :** HEK-293 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) at 1/10000 dilution

Predicted band size: 72 kDa

Observed band size: 72 kDa

Lanes 1-4: Merged signal (red and green). Green - ab199719 observed at 72 kDa. Red - loading control **ab8245** observed at 36 kDa.

ab199719 Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal was shown to specifically react with SLC27A4 / FATP4 in wild-type HEK293T cells. Loss of signal was observed when knockout cell line **ab266114** (knockout cell lysate **ab257677**) was used. Wild-type and SLC27A4 / FATP4 knockout samples were subjected to SDS-PAGE. ab199719 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HepG2 (Human liver hepatocellular carcinoma) cells labeling SLC27A4 / FATP4 with ab199719 at 1/100 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green). Cytoplasm staining on HepG2 cell line is observed. The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:-

-ve control 1: ab199719 at 1/100 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling SLC27A4 / FATP4 with ab199719 at 1/100 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/500 dilution. Cytoplasm staining on Human kidney tissue is observed. Counter stained with Hematoxylin.

Negative control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunoprecipitation - Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

SLC27A4 / FATP4 was immunoprecipitated from 1mg of HEK293 (Human embryonic kidney) whole cell lysate with ab199719 at 1/50 dilution. Western blot was performed from the immunoprecipitate using ab199719 at 1/5000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution.

Lane 1: HEK293 whole cell lysate 10µg (Input). Lane 2: HEK293 whole cell lysate following IP. Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab199719 in HEK293 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure is 3 minutes.



Western blot - Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: SLC27A4/FATP4 knockout HAP1 cell lysate (20 µg)

Lane 3: HEK293 cell lysate (20 µg)

Lane 4: HepG2 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab199719 observed at 70 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab199719 was shown to specifically react with SLC27A4/FATP4 when SLC27A4/FATP4 knockout samples were used. Wild-type and SLC27A4/FATP4 knockout samples were subjected to SDS-PAGE. ab199719 and **ab8245** (loading control to GAPDH) were diluted 1/2000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

All lanes : Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719) at 1/10000 dilution

Lane 1 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate

Lane 2 : HepG2 (Human liver hepatocellular carcinoma) whole cell lysate

Lane 3 : HEK293 (Human embryonic kidney) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 72 kDa

Observed band size: 72 kDa

Exposure time: 1 minute

Blocking/dilution buffer: 5% NFD/MTBST.



All lanes : Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719) at 1/2000 dilution

Lane 1 : Human fetal brain

Lane 2 : Human fetal kidney

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 72 kDa

Observed band size: 72 kDa

Exposure time: 1 minute

Blocking/dilution buffer: 5% NFD/MTBST.

Why choose a recombinant antibody?

<p>Research with confidence Consistent and reproducible results</p>	<p>Long-term and scalable supply Recombinant technology</p>
<p>Success from the first experiment Confirmed specificity</p>	<p>Ethical standards compliant Animal-free production</p>

Anti-SLC27A4 / FATP4 antibody [EPR17319] - C-terminal (ab199719)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors