

Anti-StAR antibody ab96637

★★★★★ [3 Abreviews](#) [21 References](#) [画像数 5](#)

製品の概要

製品名	Anti-StAR antibody
製品の詳細	Rabbit polyclonal to StAR
由来種	Rabbit
アプリケーション	適用あり: ICC, WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Cow, Dog, Pig 
免疫原	Recombinant protein fragment corresponding to a region within amino acids 15 - 238 of Human StAR (NP_000340).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.00 Preservative: 0.025% Proclin 300 Constituents: 79% PBS, 20% Glycerol (glycerin, glycerine)
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab96637の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC		Use at an assay dependent concentration.
WB	★★★★★ (3)	1/100 - 1/10000. Predicted molecular weight: 32 kDa.
IHC-P		1/100 - 1/1000.

ターゲット情報

機能	Plays a key role in steroid hormone synthesis by enhancing the metabolism of cholesterol into pregnenolone. Mediates the transfer of cholesterol from the outer mitochondrial membrane to the inner mitochondrial membrane where it is cleaved to pregnenolone.
組織特異性	Expressed in gonads, adrenal cortex and kidney.
パスウェイ	Steroid metabolism; cholesterol metabolism.
関連疾患	Defects in STAR are the cause of adrenal hyperplasia type 1 (AH1) [MIM:201710]. The most severe form of adrenal hyperplasia. It is a condition characterized by onset of profound adrenocortical insufficiency shortly after birth, hyperpigmentation reflecting increased production of pro-opiomelanocortin, elevated plasma renin activity as a consequence of reduced aldosterone synthesis, and male pseudohermaphroditism resulting from deficient fetal testicular testosterone synthesis. Affected individuals are phenotypic females irrespective of gonadal sex, and frequently die in infancy if mineralocorticoid and glucocorticoid replacement are not instituted.
配列類似性	Contains 1 START domain.
細胞内局在	Mitochondrion.

画像




Anti-StAR antibody (ab96637) at 1/1000 dilution + K562 (human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate at 30 µg

Secondary

HRP-conjugated anti-rabbit IgG antibody

Predicted band size: 32 kDa


12% SDS-PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-StAR antibody (ab96637)


Immunohistochemical analysis of paraffin-embedded human hepatoma tissue staining StAR protein at cytosol with ab96637 at 1/500.

Antigen Retrieval: EDTA based buffer, pH 8.0, 15min.



Immunocytochemistry - Anti-StAR antibody (ab96637)

Immunofluorescence analysis of 4% paraformaldehyde-fixed HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate labeling StAR protein at mitochondria with ab96637 at 1/500 dilution. Blue: Hoechst 33342 staining.



Western blot - Anti-StAR antibody (ab96637)

All lanes : Anti-StAR antibody (ab96637) at 1/5000 dilution

Lane 1 : Non-transfected HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

Lane 2 : Transfected HEK-293T whole cell lysate


Lysates/proteins at 30 μg per lane.

Secondary

All lanes : HRP-conjugated anti-rabbit IgG antibody

Predicted band size: 32 kDa

12% SDS-PAGE



All lanes : Anti-StAR antibody (ab96637) at 1/1000 dilution

Lane 1 : Whole tissue lysate prepared from murine adrenal gland

Lane 2 : Whole tissue lysate prepared from rat adrenal gland

Lysates/proteins at 50 µg per lane.

Secondary

All lanes : HRP conjugated goat anti-rabbit polyclonal at 1/2500 dilution

Developed using the ECL technique.

Predicted band size: 32 kDa

Exposure time: 20 seconds

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors