abcam

Product datasheet

Anti-STAT5a (phospho S726) antibody [EPR1914(2)] - BSA and Azide free ab248206

יילעבער RabMAb

画像数 4

製品の概要

製品名 Anti-STAT5a (phospho S726) antibody [EPR1914(2)] - BSA and Azide free

製品の詳細 Rabbit monoclonal [EPR1914(2)] to STAT5a (phospho S726) - BSA and Azide free

由来種 Rabbit

アプリケーション 適用あり: Dot blot, WB

適用なし: FuncS,ICC/IF or IHC-P

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

特記事項 ab248206 is the carrier-free version of ab128896.

> Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cellbased assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

1

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C. Do Not Freeze.

バッファー pH: 7.2

Constituent: PBS

キャリア・フリー はい

精製度 Protein A purified

ポリモノ モノクローナル **ウローン名** EPR1914(2)

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab248206の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Dot blot		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 90 kDa (predicted molecular weight: 90 kDa).

追加情報 Is unsuitable for FuncS,ICC/IF or IHC-P.

ターゲット情報

機能 Carries out a dual function: signal transduction and activation of transcription. Binds to the GAS

element and activates PRL-induced transcription.

配列類似性 Belongs to the transcription factor STAT family.

Contains 1 SH2 domain.


翻訳後修飾 Tyrosine phosphorylated in response to IL-2, IL-3, IL-7, IL-15, GM-CSF, growth hormone,

prolactin, erythropoietin and thrombopoietin. Tyrosine phosphorylation is required for DNA-binding activity and dimerization. Serine phosphorylation is also required for maximal

transcriptional activity.


細胞内局在 Cytoplasm. Nucleus. Translocated into the nucleus in response to phosphorylation.

画像



Dot Blot - Anti-STAT5a (phospho S726) antibody [EPR1914(2)] - BSA and Azide free (ab248206)

This data was developed using **ab128896**, the same antibody clone in a different buffer formulation.Primary antibody dilution: 1/1000 Secondary antibody: goat anti-rabbit lgG, (H+L), peroxidase conjugated Secondary antibody dilution: 1/2500 Blocking & dilution buffer: 5% NFDM/TBST Lane 1 sample: STAT5a (pS726) phospho peptide Lane 2 sample: STAT5a non-phospho peptide Exposure time: 10 seconds



Western blot - Anti-STAT5a (phospho S726) antibody [EPR1914(2)] - BSA and Azide free (ab248206)

All lanes : Anti-STAT5a (phospho S726) antibody [EPR1914(2)] (ab128896) at 1/5000 dilution

Lane 1 : SH-SY5Y (human neuroblastoma) treated with Okadaic acid and Calyculin A whole cell lysates

Lane 2 : Untreated SH-SY5Y (human neuroblastoma) whole cell lysates

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat anti-rabbit lgG, (H+L), peroxidase conjugated at 1/1000 dilution

Predicted band size: 90 kDa **Observed band size:** 90 kDa

Exposure time: 3 minutes

This data was developed using <u>ab128896</u>, the same antibody clone in a different buffer formulation.

Blocking and dilution buffer: 5% NFDM/TBST



(ab248206)

All lanes : Anti-STAT5a (phospho S726) antibody [EPR1914(2)] (ab128896) at 1/1000 dilution

Lane 1: SH-SY5Y cell lysates: untreated

Lane 2: SH-SY5Y cell lysates: treated with Okadaic acid and

Calyculin A

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 90 kDa Observed band size: 90 kDa

This data was developed using <u>ab128896</u>, the same antibody clone in a different buffer formulation.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors