

Anti-Stathmin 1 (phospho S63) antibody [EPR1574] ab76583

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **6 References** 画像数 7

製品の概要

製品名	Anti-Stathmin 1 (phospho S63) antibody [EPR1574]
製品の詳細	Rabbit monoclonal [EPR1574] to Stathmin 1 (phospho S63)
由来種	Rabbit
特異性	This antibody only detects Stathmin 1 phosphorylated on Serine 62. The antibody immunogen shares 86% homology with Stathmin-2, therefore it is possible that the antibody will cross-react with Stathmin-2 when phosphorylated at serine 97. This has not been assessed experimentally.
アプリケーション	適用あり: Dot blot, IHC-P, WB, IP 適用なし: Flow Cyt or ICC/IF
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HeLa cell lysates treated with Calyculin A; IHC-P: human brain tissue. IP: HeLa.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA</p>

精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR1574
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab76583の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
Dot blot		1/1000.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB	★★★★★ (1)	1/2500 - 1/10000. Predicted molecular weight: 17 kDa.
IP		1/20.

追加情報 Is unsuitable for Flow Cyt or ICC/IF.

ターゲット情報

機能	Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser-16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear.
組織特異性	Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients with chronic lymphoid or myeloid leukemia.
配列類似性	Belongs to the stathmin family. Contains 1 SLD (stathmin-like) domain.
翻訳後修飾	Many different phosphorylated forms are observed depending on specific combinations among the sites which can be phosphorylated. MAPK is responsible for the phosphorylation of stathmin in response to NGF. Phosphorylation at Ser-16 seems to be required for neuron polarization (By similarity). Phosphorylation at Ser-63 reduces tubulin binding 10-fold and suppresses the MT polymerization inhibition activity.
細胞内局在	Cytoplasm > cytoskeleton.

画像



All lanes : Anti-Stathmin 1 (phospho S63) antibody [EPR1574] (ab76583) at 1/1000 dilution

Lane 1 : Untreated HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysates

Lane 2 : HeLa (Human epithelial cell line from cervix adenocarcinoma) treated with nocodazole at 100 ng/mL for 18 hours. Whole cell lysates

Lane 3 : HeLa (Human epithelial cell line from cervix adenocarcinoma) treated with nocodazole at 100 ng/mL for 18 hours. Whole cell lysates. Then the membrane was incubated with phosphatase.

Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 17 kDa

Observed band size: 17 kDa


Exposure time: 30 seconds

Blocking/Diluting buffer and concentration 2% BSA/TBST



ab76583, at a 1/250 dilution, staining Stathmin 1 in paraffin embedded human brain tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunoprecipitation - Anti-Stathmin 1 (phospho S63) antibody [EPR1574] (ab76583)

Stathmin 1 was immunoprecipitated from 0.35 mg HeLa (Human cervix adenocarcinoma epithelial cell) treated with Calyculin A whole cell lysate 10 µg with ab76583 at 1/50 dilution (2µg). VeriBlot for IP Detection Reagent (HRP)([ab131366](#)) was used at 1/5000 dilution.


Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) treated with Calyculin A whole cell lysate 10 µg

Lane 2: ab76583 IP in HeLa treated with Calyculin A whole cell lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab76583 in HeLa treated with Calyculin A whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

75kDa band could be stathmin/alpha tubulin complex.
(PMID:9369201)



Western blot - Anti-Stathmin 1 (phospho S63) antibody [EPR1574] (ab76583)

All lanes : Anti-Stathmin 1 (phospho S63) antibody [EPR1574] (ab76583) at 1/10000 dilution

Lane 1 : HeLa cell lysate, untreated

Lane 2 : HeLa cell lysate treated with Calyculin A


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled goat anti rabbit at 1/2000 dilution

Predicted band size: 17 kDa


Observed band size: 17 kDa



Dot blot analysis of Stathmin 1 (phospho S63) phospho peptide (Lane 1) and Stathmin 1 non-phospho peptide (Lane 2) labeling Stathmin 1 (phospho S63) with ab76583 at a dilution of 1/1000. [ab97051](#) (Peroxidase conjugated goat anti-rabbit IgG) (H+L) at 1/100 000 was used as the secondary antibody.

Blocking and diluting buffer: 5% NFDm/TBST.

Exposure time: 3 minutes.



Alkaline phosphatase treatment removes S63 phosphorylation. Samples treated with phosphatase were run alongside the normal lysate, and the phosph-S63 signal is not detected after phosphatase treatment, thus suggesting the signal is very specific to the phosphorylated S63.

Whole cell lysate prepared from a human colon cancer cell line was loaded at 20µg.

ab76583 used at a 1/1000 dilution.

Secondary used was an HRP conjugated goat anti-rabbit polyclonal used at a 1/10000 dilution.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Stathmin 1 (phospho S63) antibody [EPR1574]
(ab76583)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors