

### Anti-STING antibody [SP338] ab227704

リコンビナント **RabMAb**

**2 References**   [画像数 19](#)

#### 製品の概要

製品名	Anti-STING antibody [SP338]
製品の詳細	Rabbit monoclonal [SP338] to STING
由来種	Rabbit
アプリケーション	<b>適用あり:</b> Flow Cyt (Intra), IHC-P, WB, ICC/IF
種交差性	<b>交差種:</b> Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Human lymph node, thymus, tonsil, colon, spleen, colon adenocarcinoma, lung, lung squamous cell carcinoma, lung adenocarcinoma and pancreatic adenocarcinoma tissues. WB: THP-1 cell lysate. Flow Cyt: THP-1 cells. ICC/IF: THP-1 cells.
特記事項	<b>This product is FOR RESEARCH USE ONLY. For commercial use, please contact <a href="mailto:partnerships@abcam.com">partnerships@abcam.com</a>.</b>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.60 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
精製度	Protein A/G purified
特記事項(精製)	Purified from TCS by protein A/G.
ポリ/モノ	モノクローナル
クローン名	SP338
アイソタイプ	IgG

#### アプリケーション

## The Abpromise guarantee

**Abpromise保証は、**次のテスト済みアプリケーションにおけるab227704の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/20 - 1/400.
IHC-P		1/100. Perform heat mediated antigen retrieval with EDTA buffer pH 8.0 before commencing with IHC staining protocol.
WB		1/400. Predicted molecular weight: 42 kDa.
ICC/IF		1/50.

## ターゲット情報


機能	Facilitator of innate immune signaling that promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm. Able to activate both NF-kappa-B and IRF3 transcription pathways to induce expression of type I interferon and exert a potent anti-viral state following expression. May be involved in translocon function, the translocon possibly being able to influence the induction of type I interferons. May be involved in transduction of apoptotic signals via its association with the major histocompatibility complex class II (MHC-II). Mediates death signaling via activation of the extracellular signal-regulated kinase (ERK) pathway.
組織特異性	Ubiquitously expressed.
配列類似性	Belongs to the TMEM173 family.
翻訳後修飾	Phosphorylated on tyrosine residues upon MHC-II aggregation (By similarity). Phosphorylated on Ser-358 by TBK1, leading to activation and production of IFN-beta. Ubiquitinated. 'Lys-63'-linked ubiquitination mediated by TRIM56 at Lys-150 promotes homodimerization and recruitment of the antiviral kinase TBK1 and subsequent production of IFN-beta. 'Lys-48'-linked polyubiquitination at Lys-150 occurring after viral infection is mediated by RNF5 and leads to proteasomal degradation.
細胞内局在	Endoplasmic reticulum membrane. Mitochondrion outer membrane. Cell membrane. Cytoplasm > perinuclear region. In response to double-stranded DNA stimulation, relocalizes to perinuclear region, where the kinase TBK1 is recruited.

## 画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat spleen tissue sections labeling TMEM173 with ab227704 at 1/100 dilution (1.05 µg/ml). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10mins. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Hematoxylin was used as a counterstain. Positive staining on the rat spleen, performed on a Leica Biosystems BOND™ RX instrument. The section was incubated with ab227704 for 10 mins at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human colon tissue sections labeling TMEM173 with ab227704 at 1/100 dilution (1.05 µg/ml). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10mins. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Hematoxylin was used as a counterstain. Positive staining on the human colon, performed on a Leica Biosystems BOND™ RX instrument. The section was incubated with ab227704 for 10 mins at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human tonsil tissue sections labeling TMEM173 with ab227704 at 1:100 dilution (1.05 µg/ml). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10mins. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Hematoxylin was used as a counterstain. Positive staining on the human tonsil, performed on a Leica Biosystems BOND™ RX instrument.


The section was incubated with ab227704 for 10 mins at room temperature.



Western blot - Anti-STING antibody [SP338] (ab227704)


Anti-STING antibody [SP338] (ab227704) at 1/400 dilution + THP-1 (human monocytic leukemia cell line) cell lysate

**Predicted band size: 42 kDa**



Immunocytochemistry/ Immunofluorescence - Anti-STING antibody [SP338] (ab227704)


Immunocytochemistry/ Immunofluorescence analysis of THP-1 (human monocytic leukemia monocyte) cells labeling TMEM173 with purified ab227704 at 1/50 (2.1 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with **ab195889** Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-STING antibody [SP338] (ab227704)


Flow cytometry analysis of THP-1 (human monocytic leukemia monocyte) labeling TMEM173 with purified ab227704 at 1/20 dilution (5.25 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, **ab150081**) at 1/2000 dilution was used as a secondary antibody. Isotype control -Rabbit monoclonal IgG (**ab172730**) / Black. Unlabeled control -Unlabelled cells / blue.






Formalin-fixed, paraffin-embedded human lymph node tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Formalin-fixed, paraffin-embedded human thymus tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Formalin-fixed, paraffin-embedded human tonsil tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Flow cytometric analysis of THP-1 (human monocytic leukemia cell line) cell line labeling TMEM173 with ab227704 at 1/400 dilution (green) compared with a rabbit IgG negative control (blue).

Flow Cytometry (Intracellular) - Anti-STING antibody [SP338] (ab227704)



Formalin-fixed, paraffin-embedded human colon tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)



Formalin-fixed, paraffin-embedded human spleen tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)






Formalin-fixed, paraffin-embedded human colon adenocarcinoma tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Formalin-fixed, paraffin-embedded human lung tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)




Formalin-fixed, paraffin-embedded human lung adenocarcinoma tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)



Formalin-fixed, paraffin-embedded human pancreatic adenocarcinoma tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)



Formalin-fixed, paraffin-embedded human pancreatic adenocarcinoma tissue stained for TMEM173 using ab227704 at 1/100 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STING antibody [SP338] (ab227704)

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-STING antibody [SP338] (ab227704)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors