

Anti-Synaptophysin antibody [SP11] ab16659

RabMAb

★★★★★ **2 Abreviews** **16 References** 画像数 5

製品の概要

製品名	Anti-Synaptophysin antibody [SP11]
製品の詳細	Rabbit monoclonal [SP11] to Synaptophysin
由来種	Rabbit
アプリケーション	適用あり: ICC, WB, IHC-P, IHC-Fr
種交差性	交差種: Rat, Human
免疫原	Synthetic peptide within Human Synaptophysin aa 250 to the C-terminus. The exact sequence is proprietary. Database link: P08247
エピトープ	C-terminus
ポジティブ・コントロール	WB: Human Brain lysate. IHC-Fr: Human Pancreas tissue sections. ICC: Primary hippocampal rat neurons/glia cells. IHC: human neuroendocrine tumor and Human pancreas.
特記事項	This product is FOR RESEARCH USE ONLY. For commercial use, please contact partnerships@abcam.com.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.50 Preservative: 0.1% Sodium azide Constituents: Tissue culture supernatant, Tris buffered saline, 1% BSA

精製度	Tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	SP11
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab16659の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC		Use a concentration of 5 µg/ml.
WB		1/200. Predicted molecular weight: 38 kDa.
IHC-P		1/200. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
IHC-Fr		1/200.

ターゲット情報

機能	Possibly involved in structural functions as organizing other membrane components or in targeting the vesicles to the plasma membrane. Involved in the regulation of short-term and long-term synaptic plasticity.
組織特異性	Characteristic of a type of small (30-80 nm) neurosecretory vesicles, including presynaptic vesicles, but also vesicles of various neuroendocrine cells of both neuronal and epithelial phenotype.
関連疾患	Mental retardation, X-linked, SYP-related
配列類似性	Belongs to the synaptophysin/synaptobrevin family. Contains 1 MARVEL domain.
ドメイン	The calcium-binding activity is thought to be localized in the cytoplasmic tail of the protein.
翻訳後修飾	Ubiquitinated; mediated by SIAH1 or SIAH2 and leading to its subsequent proteasomal degradation.
細胞内局在	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Cell junction, synapse, synaptosome.

画像




Immunocytochemistry - Anti-Synaptophysin antibody [SP11] (ab16659)

ab16659 staining Synaptophysin in primary hippocampal rat neurons/glia, (obtained from Neuromics, cat. no. PC35101), DIV14. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab16659 at 1µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).


Also suitable in cells fixed with 4% paraformaldehyde (10 min).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Synaptophysin antibody [SP11] (ab16659)

Immunohistochemical staining of human neuroendocrine tumor with ab16659.



Immunohistochemical staining of human pancreas with ab16659.


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Synaptophysin antibody [SP11] (ab16659)



IHC image of Synaptophysin staining in a section of frozen normal human pancreas performed on a Leica BOND™ system using the standard protocol. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with ab16659, 1/200 dilution, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

Immunohistochemistry (Frozen sections) - Anti-Synaptophysin antibody [SP11] (ab16659)

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Anti-Synaptophysin antibody [SP11] (ab16659) at 1/500 dilution +
Lysate prepared from human brain

Predicted band size: 38 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors