

Anti-Talin 1 antibody [93E12] ab104913

5 References

製品の概要

製品名	Anti-Talin 1 antibody [93E12]
製品の詳細	Mouse monoclonal [93E12] to Talin 1
由来種	Mouse
特異性	ab104913 can be used for specific detection of the focal adhesion protein Talin 1 and does not cross react with Talin 2.
アプリケーション	適用あり: ELISA, WB, ICC/IF, IHC-P, IP
種交差性	交差種: Human
免疫原	Recombinant fragment, corresponding to amino acids 780-850 of Talin 1.
ポジティブ・コントロール	Fibroblasts
特記事項	<p>Growth Conditions: RPMI1650 + Glu + Pen + Strep + 10% FCS @ 37C + 5%CO₂.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	<p>Preservative: 0.02% Sodium azide</p> <p>Constituent: 99.98% PBS</p>
精製度	Protein A purified
一次抗体 備考	Growth Conditions: RPMI1650 + Glu + Pen + Strep + 10% FCS @ 37C + 5%CO ₂ .
ポリ/モノ	モノクローナル

クローン名	93E12
ミエローマ	NS0/1
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab104913の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

ターゲット情報

機能	Probably involved in connections of major cytoskeletal structures to the plasma membrane. High molecular weight cytoskeletal protein concentrated at regions of cell-substratum contact and, in lymphocytes, at cell-cell contacts.
配列類似性	Contains 1 FERM domain. Contains 1 ILWEQ domain.
細胞内局在	Cell projection, ruffle membrane. Cytoplasm, cytoskeleton. Cell surface. Cell junction, focal adhesion. Colocalizes with LAYN at the membrane ruffles. Localized preferentially in focal adhesions than fibrillar adhesions (By similarity).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors